

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
10 October 2002 (10.10.2002)


PCT

(10) International Publication Number  
WO 02/078603 A2

- (51) International Patent Classification<sup>7</sup>: A61K (74) Agent: ALTMAN, Daniel, E.; KNOBBE, MARTENS, OLSON AND BEAR, LLP, 620 Newport Center Drive, 16th Floor, Newport Beach, CA 92660 (US).
- (21) International Application Number: PCT/US02/10071
- (22) International Filing Date: 29 March 2002 (29.03.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/281,130 2 April 2001 (02.04.2001) US
- (71) Applicant (for all designated States except US): THE GOVERNMENT OF THE UNITED STATES OF AMERICA, as represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; National Institutes of Health, Office of Technology Transfer, 6011 Executive Blvd., Suite 325, Rockville, MD 20852-3804 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): MAYER, Ghislaine [HT/US]; 19902 Sweetgum Circle #43, Germantown, MD 20874 (US). MILLER, Louis, H. [US/US]; 5450 Whitley Park Terrace #609, Bethesda, MD 20814 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EC, EE, EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Declaration under Rule 4.17:  
— of inventorship (Rule 4.17(iv)) for US only

[Continued on next page]

(54) Title: PLASMODIUM FALCIPARUM ERYTHROCYTE BINDING PROTEIN BAEBL FOR USE AS A VACCINE

(57) Abstract: The invention relates to *lasmodium falciparum* Erythrocyte Binding Protein BAEBL for use as a vaccine.

BEST AVAILABLE COPY

WO 02/078603 A2



**Published:**

— without international search report and to be republished  
upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**PLASMODIUM FALCIPARUM ERYTHROCYTE BINDING PROTEIN BAEBL  
FOR USE AS A VACCINE**

5

**Field of the Invention**

The invention relates to *Plasmodium falciparum* Erythrocyte Binding Protein BAEBL for use as a vaccine.

**Background of the Invention**

10 The erythrocytic stage of *Plasmodium falciparum* causes several million deaths yearly, primarily in Africa. The parasite lives within the erythrocyte except during the brief period when merozoites, the invasive stage of the parasite, are released from infected erythrocytes to invade uninfected erythrocytes. Invasion of erythrocytes by merozoites is a multistep process that includes: attachment, reorientation of the merozoite in such a way that its apical end is in contact with the erythrocyte surface, junction formation, and entry  
15 into the parasitophorous vacuole (Dvorak, J.A. *et al.* 1975 *Science* 187:748-9; Aikawa, M., *et al.* 1978 *J Cell Biol* 77:72-82). The binding of merozoites to erythrocytes requires parasite receptors (Camus, D. & Hadley, T.H. 1985 *Science* 230:553-556; Haynes, J.D. *et al.* 1988 *J Exp Med* 167:1873-1881; Adams, J.H. *et al.* 1990 *Cell* 63:142-153; Sim, B.K.L., *et al.* 1990 *J Cell Biol* 111:1877-1884; Galinski, M.R. *et al.* 1992 *Cell* 69:1213-1226).

20 One family of these parasite receptors is named Duffy binding-like erythrocyte binding protein (DBL-EBP) for its similarity to the *P. vivax* and *P. knowlesi* proteins that bind to the Duffy blood group antigens (Duffy positive) on human erythrocytes (Adams, J.H. *et al.* 1992 *PNAS USA* 89:7085-7089). *P. vivax* does not infect Africans lacking the Duffy blood group antigens (Duffy negative), and *P. knowlesi* will not form a junction with  
25 or invade Duffy negative human erythrocytes (Miller, L.H. *et al.* 1976 *N Engl J Med* 295:302-304; Miller, L.H., *et al.* 1979 *J Exp Med* 149:172-184). Region II, a domain of the *P. vivax* DBL-EBP, has the same specificity as the full-length protein (Chitnis, C.E. & Miller, L.H. 1994 *J Exp Med* 180:497-506).

*P. knowlesi* has three highly homologous DBL proteins, each with different  
30 specificities as defined by region II (Haynes, J.D. *et al.* 1988 *J Exp Med* 167:1873-1881; Ranjan, A. & Chitnis, C.E. 1999 *PNAS USA* 96:14067-14072). One binds to Duffy blood group antigens on human and rhesus erythrocytes, a second binds to sialic acid on rhesus

erythrocytes, and a third binds to an unidentified receptor on rhesus erythrocytes. Whereas *P. knowlesi* can only invade Duffy positive human erythrocytes, it can invade rhesus erythrocytes that have been rendered Duffy negative by protease treatment and by removal of sialic acid with neuraminidase (Haynes, J.D. *et al.* 1988 *J Exp Med* 167:1873-1881; Miller, L.H. *et al.* 1973 *J Exp Med* 138:1597-1601). *P. knowlesi* invades these enzymatically treated erythrocytes at the same rate as the untreated erythrocytes, indicating a highly efficient alternative pathway of invasion.

The Duffy binding proteins of *P. vivax* (PvDBP) and *P. knowlesi* (PkDBP) are part of a larger family of *Plasmodium* proteins that include EBA-175 of *P. falciparum*. EBA-175 binds to sialic acid and the peptide backbone of glycophorin A on the erythrocyte surface (Sim, B.K.L. *et al.* 1994 *Science* 264:1941-1944). As in the case of *P. vivax*, the binding domain of EBA-175 is defined by region II. Unlike *P. vivax*, which cannot infect Duffy negative erythrocytes, some strains of *P. falciparum* parasites have alternative pathways of invasion, not requiring glycophorin A for either invasion or growth *in vitro*. Thus, other receptors must be involved in these alternative pathways (Dolan, S.A. *et al.* 1990 *J Clin Invest* 86:618-624).

The *P. falciparum* genome sequence identifies at least four paralogues of EBA-175. We have studied one of these DBL genes of *P. falciparum*, named *baebl* (Adams, J.H. *et al.* 2001 *Trends Parasitol* 17:297-9), to explore its possible role in invasion.

### Summary of the Invention

The present invention relates to *Plasmodium falciparum* Erythrocyte Binding Protein BAEBL for use as a vaccine. A BAEBL polynucleotide sequence or a portion thereof, or a BAEBL polypeptide sequence or a portion thereof, is used to induce an immune response to a Plasmodium parasite, whereby a human is protected against malaria. The BAEBL polynucleotide sequence is alternatively used to express recombinant polypeptides or portions thereof. Furthermore, synthetic BAEBL polypeptides or portions thereof are prepared in accordance with the present invention.

### Brief Description of the Drawings

**Figure 1.** Sequencing strategy and exon/intron structure of *baebl*. Oligonucleotides were designed based on the genomic sequence obtained from the *Plasmodium falciparum* genome project (Sanger Centre) and used for sequencing of

genomic DNA (GenBank No. AF332918) and RT-PCR of mRNA (GenBank No. AF332919) to determine the intron/exon structure in *P. falciparum* Dd2/Nm strain. (A) Schematic representation of the gene and predicted protein structure of *baebl*. Predicted protein structure has strong similarity with EBA-175, containing the putative signal sequence (SS; aa 1-21) predicted by SIGNALP V2.0; region II (two Duffy binding-like (DBL) domains, F1 and F2); region VI (3' Cys), the transmembrane domain (TM; aa 1134-1153) predicted by TMHMM V2.0, followed by the putative cytoplasmic domain (Cyt). (B) f1 to f9 primers (see Examples) are used for RT-PCR of mRNA (lanes marked c) and PCR of genomic DNA (lanes marked g). Bar = 1 kb.

**Figure 2.** Confocal microscopy demonstrates the localization of BAEBL in micronemes. (A) Dd2/Nm schizonts were double labeled with anti-BAEBL region II and anti-EBA-175. Schizonts immunolabeled with anti-BAEBL region II were stained with Alexa 488 secondary antibody. Schizonts labeled with anti-EBA-175 were stained with Alexa 594 secondary antibody. (B) Dd2/Nm schizonts were double labeled with anti-BAEBL region VI and anti-EBA-175. Schizonts immunolabeled anti-BAEBL region VI were stained with Alexa 488 secondary antibody. Schizonts labeled with anti-EBA-175 were stained with Alexa 594 secondary antibody. (C) Dd2/Nm schizonts were double labeled with anti-BAEBL region II and anti-RAP-1 monoclonal antibody. Schizonts immunolabeled anti-BAEBL region II were stained with Alexa 488 secondary antibody. Schizonts labeled with anti-RAP-1 were stained with TRITC secondary antibody. (D) Dd2/Nm schizonts were double labeled with anti-BAEBL region VI and anti-RAP-1 monoclonal antibody. Schizonts immunolabeled with anti-BAEBL region VI were stained with Alexa 488 secondary antibody. Schizonts labeled with anti-RAP-1 were stained with TRITC secondary antibody.

**Figure 3.** Evidence that anti-region II (Anti-R2) and anti-region VI (Anti-R6) sera immunoprecipitate the same protein. The supernatant was preabsorbed with either anti-region 2 or anti-region 6 followed by immunoprecipitation by the two sera. BAEBL was removed by both sera.

**Figure 4.** BAEBL and EBA-175 did not bind to neuraminidase (NM eluate) or trypsin-treated erythrocytes (Trypsin RBC). Eluates of BAEBL and EBA-175 were only seen from normal erythrocytes.

**Figure 5.** BAEBL binds and is eluted from En(a-) erythrocytes that lack glycophorin A. NM RBC are neuraminidase-treated normal erythrocytes.

**Figure 6.** Absorption and elution of BAEBL (A, B) and EBA-175 (C, D) with various amounts (25, 50, 2 × 50 µl, and 4 × 50 µl of packed erythrocytes) of Gerbich [Ge(-2, -3, 4)], normal, and neuraminidase (NM)-treated erythrocytes. For elution, 25 µl and 50 µl of packed erythrocytes were used.

#### Summary of Sequences

SEQ ID NO: 1 and Figure 7 are the genomic sequence encoding *Plasmodium falciparum* Erythrocyte Binding Protein BAEBL; start and stop codons are indicated in bold, and the introns span nucleotides 3499-3638, 3718-3846, and 3930-4061.

SEQ ID NO: 2 is the amino acid sequence encoding *Plasmodium falciparum* Erythrocyte Binding Protein BAEBL.

#### Detailed Description of the Preferred Embodiment

A member of a *Plasmodium* receptor family for erythrocyte invasion was identified on chromosome 13 from the *Plasmodium falciparum* genome sequence of the Sanger Centre. The protein (named BAEBL) has homology to EBA-175, a *P. falciparum* receptor that binds specifically to sialic acid and the peptide backbone of glycophorin A on erythrocytes. Both EBA-175 and BAEBL localize to the micronemes, an organelle at the invasive end of the parasite that contains other members of the family. Like EBA-175, the erythrocyte receptor for BAEBL is destroyed by neuraminidase and trypsin, indicating that the erythrocyte receptor is a sialoglycoprotein. Its specificity, however, differs from EBA-175 in that BAEBL can bind to erythrocytes that lack glycophorin A, the receptor for EBA-175. It has reduced binding to erythrocytes with the Gerbich mutation found in another erythrocyte sialoglycoprotein (glycophorin C/D). The interest in BAEBL's reduced binding to Gerbich erythrocytes derives from the high frequency of the Gerbich phenotype in some regions of Papua New Guinea where *P. falciparum* is hyperendemic.

The present invention relates, in general, to a substantially purified polynucleotide sequence (e.g. a DNA sequence) encoding all, or a portion, of BAEBL of the DBL-EBP family of a *Plasmodium* parasite (particularly, *Plasmodium falciparum*). A "portion" as used herein preferably consists of at least five (or six) amino acids or, correspondingly, at

least 15 (or 18) nucleotides. GenBank No. AF332918 encodes the genomic DNA and GenBank No. AF332919 encodes the cDNA of BAEBL.

5 The present invention further relates to a polynucleotide sequence encoding a BAEBL protein of other *Plasmodium* parasites such as, for example, *P. vivax* or *P. knowlesi*. One of ordinary skill in the art, given the present disclosure, could easily identify and clone analogous genes in such species without undue experimentation.

10 In one embodiment, the present invention relates to a polynucleotide sequence given in SEQ ID NO: 1 encoding the entire amino acid sequence of BAEBL (the specific DNA sequence defined therein being only an example). The polynucleotide sequence can be genomic DNA or cDNA. Polynucleotide sequences to which this invention relates also include those encoding substantially the same protein as that encoded by SEQ ID NO: 1, which include, for example, allelic forms of the given amino acid sequences and alternatively spliced products.

15 The present invention relates to a recombinant DNA molecule comprising a vector and a DNA sequence encoding BAEBL, or a portion thereof. Using methodology well known in the art, recombinant DNA molecules of the present invention can be constructed. Possible vectors for use in the present invention include expression vectors. The *Plasmodium* BAEBL encoding sequences of the present invention can be inserted into commercially available DNA vectors (expression vectors) to express the encoded protein product. The expression vectors have promoter sequences and other regulatory sequences necessary for expression in host cells. The technique of using expression vectors to introduce exogenous genes and express their protein products in a host cell is well known to those familiar with the art. For example the expression vector pET21a is commercially available and can be used to express proteins in *E. coli*. Alternatively the protein can be expressed in a eukaryotic cell, such as yeast, using *Pichia* expression vectors (i.e. pHIL-D2) commercially available from Invitrogen. The baculovirus system is also commercially available and can be used to express the BAEBL genes in insect cultures.

25 Once the *baebl* gene or fragment thereof has been cloned into an expression vector, the resulting vector can be used to transform a host cell, using procedures known to those familiar with the art. Such transformation procedures include but are not limited to microinjection, microprojectile bombardment, electroporation, calcium chloride

30

permeabilization, polyethylene glycol permeabilization, protoplast fusion or bacterial mediated mechanisms such as *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes*.

Host cells may be selected from any cell in which expression of modified proteins can be made compatible, including bacteria, fungus, yeast, plant cells and animal cells.

5 Suitable host cells include prokaryotes selected from the genus *Escherichia* or *Staphylococcus* and eukaryotes selected from the genus *Pichia* (including *Saccharomyces cerevisiae*, for example). In addition, mammalian cell culture (such as CHO and COS cells) can be used to express the BAEBL proteins and peptide fragments.

The transformed host cells synthesize the BAEBL protein or peptide fragment

10 which can be isolated and purified using standard methods known to those familiar with the art. In one embodiment the BAEBL proteins and peptide fragments can be expressed as fusion proteins to assist in the purification of the BAEBL protein products.

The present invention also relates to a *Plasmodium* BAEBL protein separated from those proteins with which it is naturally associated. One skilled in the art can easily purify

15 BAEBL using methodologies well known in the art.

The present invention further relates to a recombinantly produced BAEBL protein with the amino acid sequence given in SEQ ID NO: 2, an allelic variation thereof or a chimeric protein thereof. The present invention also relates to recombinantly produced peptide fragments of BAEBL. Further, the present invention relates to synthetic BAEBL or

20 a synthetic peptide fragment thereof.

The present invention further relates to a polypeptide comprising an amino acid sequence having a consecutive number of amino acid sequences selected from a BAEBL protein, for example, *Plasmodium* BAEBL protein having the sequence of SEQ ID NO: 2, which are useful as diagnostic agents or can be utilized as therapeutic agents for treating or

25 preventing malaria. Some agents are useful as antigenic fragments to display linear epitopes, and other agents are useful as antigenic fragments to display conformational epitopes for generating neutralizing antibodies. In another embodiment, the invention relates to a polypeptide comprising an amino acid sequence that encodes an EBA-175-like domain of a BAEBL protein, which is BAEBL region II, constituting two Duffy binding-like (DBL) domains, F1 and F2, for example, *Plasmodium* BAEBL protein having the

30 sequence of SEQ ID NO: 2 running from amino acids 154-738. In another embodiment, the invention is directed to a polypeptide comprising an amino acid sequence having the



following number of consecutive amino acids taken from a BAEBL protein, for example, *Plasmodium* BAEBL protein having the sequence of SEQ ID NO: 2: 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556,

557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, and 584.

The present invention further relates to antibodies specific for epitopes present on the BAEBL proteins of the *Plasmodium* parasites. Thus, in one embodiment, the present invention relates to antibodies (such as monoclonal, polyclonal, chimeric, humanized, and anti-idiotypic) specific for the BAEBL protein. More particularly, the antibodies are directed to conserved regions of BAEBL and more preferably to BAEBL region II. One skilled in the art, using standard techniques well known to those skilled in the art can raise antibodies to the proteins and peptide fragments disclosed in the present invention. These antibodies can be useful as diagnostic agents or can be utilized as therapeutic agents for treating or preventing malaria.

The present invention relates to a vaccine for use in humans against malaria. As is customary for vaccines, BAEBL or a portion thereof, can be delivered to a human in a pharmacologically acceptable vehicle. As one skilled in the art will understand, it is not necessary to use the entire protein. A portion of the protein (for example, a peptide corresponding to a conserved region of the BAEBL protein) can be conjugated to pharmacologically acceptable carriers, including *diphtheria* toxoid, *pertussis* toxoid, or *tetanus* toxoid.

Vaccines of the present invention can include effective amounts of immunological adjuvants known to enhance an immune response. Adjuvants suitable for co-administration in accordance with the present invention should be ones that are potentially safe, well tolerated and effective in people. Such immunological adjuvants include QS-21, Detox-PC, MPL-SE, MoGM-CSF, TiterMax-G, CRL-1005, GERBU, TERamide, PSC97B, Adjuver, PG-026, GSK-1, GcMAF, B-aletine, MPC-026, Adjuvax, CpG ODN, Betafectin, Alum, and MF59 (see Kim *et al.* 2000 *Vaccine*, 18:597 and references therein).

The protein or peptide fragment is present in the vaccine in an amount sufficient to induce an immune response against the antigenic portion and thus to protect against *Plasmodium* infection thereby protecting the human against malaria. Protective antibodies are usually best elicited by a series of 2-3 doses given about 2 to 3 weeks apart. The series can be repeated when concentrations of circulating antibodies in the human drops. Further, the vaccine can be used to immunize a human against other forms of malaria (that is, heterologous immunization).

Compositions comprising substantially purified polynucleotide sequences encoding a BAEBL protein or peptide fragment can be used in accordance with the present invention as vaccines. Live vector viruses are contemplated, where retroviruses, adenoviruses, or adeno-associated viruses are engineered to carry polynucleotide sequences encoding a BAEBL protein or peptide fragment. Genetic immunization is an alternative, where naked DNA encoding a BAEBL protein or peptide fragment is administered to cells and the encoded protein antigens are expressed.

The present invention further yet relates to receptor blocking therapy which disrupts the life cycle of the parasite in humans. Administering to a human antibodies of the present invention specific for the binding site of the BAEBL ligand of the present invention can prevent invasion of red blood cells by the merozoite, a necessary event in the life cycle of the Plasmodium parasite. Alternatively, the erythrocyte receptor can be administered to a human, which is glycophorin C/D (Reid and Spring, 1994 *Transfusion Medicine* 4:139). The BAEBL ligand on the merozoite will bind the circulating receptor rather than the determinate on the red blood cells. Attachment of the merozoite to the red blood cells, and hence invasion of the parasite, is prevented.

The major human malaria parasite, *P. falciparum*, has redundant or alternate receptor-ligand pathways of invasion. Therefore, an effective vaccine for blocking parasite invasion of erythrocytes by *P. falciparum* malaria will also target the redundant receptor ligand interactions that occur during the invasion process. Thus in some embodiments, the present vaccine compositions comprise a BAEBL polypeptide in combination with additional *Plasmodium* specific proteins or peptide fragments. For example, the second polypeptide may comprise an amino acid sequence that encodes a Duffy binding protein or erythrocyte binding antigen-175 (EBA-175) of a malaria Plasmodium parasite. The Duffy binding protein and EBA-175 are members of the EBL family of proteins that are utilized by Plasmodium parasites to invade erythrocytes. Thus, one vaccine composition in accordance with the present invention comprises two or more proteins (or peptide fragments) and a pharmaceutically acceptable vehicle, wherein at least one protein (or peptide fragment) is Duffy binding protein or EBA-175. EBA-175 and Duffy binding proteins of Plasmodium parasites have been described in the prior art as well as their use in preparing vaccines to prevent malaria infections. See U.S. Pat Nos. 5,198,347 and 6,120,770.

The present invention also relates to a method of vaccinating a vertebrate species, particularly a human, against a malaria *Plasmodium* parasite. The method comprises the steps of administering a vaccine composition comprising a protein or peptide fragment of BAEBL where the peptide fragment comprises at least a consecutive six amino acid sequence and a physiologically acceptable vehicle. In one embodiment the vaccine composition further comprises a second protein or peptide fragment wherein the second protein or peptide fragment comprises the Duffy binding protein or erythrocyte binding antigen-175 of an erythrocyte binding protein. The vaccine composition also can include various adjuvants known to those skilled in the art. The vaccine composition can be administered to a vertebrate species either orally or parenterally using techniques well known to those skilled in the art.

#### Nucleic Acid Molecules

As indicated herein, nucleic acid molecules of the present invention may be in the form of RNA or in the form of DNA obtained by cloning or produced synthetically. The DNA may be double-stranded or single-stranded. Single-stranded DNA or RNA may be the coding strand, also known as the sense strand, or it may be the non-coding strand, also referred to as the anti-sense strand.

By "isolated" nucleic acid molecule(s) is intended a nucleic acid molecule, DNA or RNA, which has been removed from its native environment. For example, recombinant DNA molecules contained in a vector are considered isolated for the purposes of the present invention. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the DNA molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced synthetically.

Nucleic acid molecules of the present invention include DNA molecules comprising an open reading frame (ORF) of a wild-type *baebl* gene; and DNA molecules which comprise a sequence substantially different from those described above but which, due to the degeneracy of the genetic code, still encode an ORF of a wild-type BAEBL polypeptide. Of course, the genetic code is well known in the art. Degenerate variants optimized for human codon usage are preferred.

In another aspect, the invention provides a nucleic acid molecule comprising a polynucleotide which hybridizes under stringent hybridization conditions to a portion of the polynucleotide in a nucleic acid molecule of the invention described above. By "stringent hybridization conditions" is intended overnight incubation at 42°C in a solution comprising: 50% formamide, 5 times SSC (750 mM NaCl, 75 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 times Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1 times SSC at about 65 degree C.

By a polynucleotide which hybridizes to a "portion" of a polynucleotide is intended a polynucleotide (either DNA or RNA) hybridizing to at least about 15 nucleotides (nt), and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably about 30-70 nt of the reference polynucleotide.

By a portion of a polynucleotide of "at least 20 nt in length," for example, is intended 20 or more contiguous nucleotides from the nucleotide sequence of the reference polynucleotide. Of course, a polynucleotide which hybridizes only to a complementary stretch of T (or U) residues, would not be included in a polynucleotide of the invention used to hybridize to a portion of a nucleic acid of the invention, since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly T (or U) stretch or the complement thereof (e.g., practically any double-stranded DNA clone).

As indicated herein, nucleic acid molecules of the present invention which encode a BAEBL polypeptide may include, but are not limited to those encoding the amino acid sequence of the full-length polypeptide, by itself, the coding sequence for the full-length polypeptide and additional sequences, such as those encoding a leader or secretory sequence, such as a pre-, or pro- or prepro-protein sequence, the coding sequence of the full-length polypeptide, with or without the aforementioned additional coding sequences, together with additional, non-coding sequences, including for example, but not limited to introns and non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences that play a role in transcription, mRNA processing, including splicing and polyadenylation signals, for example, ribosome binding and stability of mRNA; and additional coding sequence which codes for additional amino acids, such as those which provide additional functionalities.

The present invention further relates to variants of the nucleic acid molecules of the present invention, which encode portions, analogs or derivatives of the BAEBL protein. Variants may occur naturally, such as a natural allelic variant. By an "allelic variant" is intended one of several alternate forms of a gene occupying a given locus on a genome of an organism (*Genes II*, 1985 Lewin, B., ed., John Wiley & Sons, New York). Non-naturally occurring variants may be produced using art-known mutagenesis techniques.

Such variants include those produced by nucleotide substitutions, deletions or additions, which may involve one or more nucleotides. The variants may be altered in coding regions, non-coding regions, or both. Alterations in the coding regions may produce conservative or non-conservative amino acid substitutions, deletions or additions. Especially preferred among these are silent substitutions, additions and deletions, which do not alter the properties and activities of the BAEBL polypeptide or portions thereof. Also especially preferred in this regard are conservative substitutions.

Further embodiments of the invention include nucleic acid molecules comprising a polynucleotide having a nucleotide sequence at least 70% identical, and more preferably at least 80%, 90%, 95% or 99% identical to a nucleotide sequence encoding a polypeptide having the amino acid sequence of a wild-type BAEBL polypeptide or a nucleotide sequence complementary thereto.

By a polynucleotide having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence encoding a BAEBL polypeptide is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the BAEBL polypeptide. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular nucleic acid molecule is at least 70%, 80%, 90%, 95% or 99% identical to the reference nucleotide sequence can be determined conventionally using known computer programs such as the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, Wis. 53711). Bestfit uses the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2: 482-489 (1981), to find the best segment of homology between two sequences. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference nucleotide sequence and that gaps in homology of up to 5% of the total number of nucleotides in the reference sequence are allowed.

The present application is directed to nucleic acid molecules at least 70%, 80%, 90%, 95% or 99% identical to the nucleic acid sequences shown herein in the Sequence Listing which encode a polypeptide having BAEBL polypeptide activity. By "a polypeptide having BAEBL activity" is intended polypeptides exhibiting BAEBL activity in a particular biological assay. For example, BAEBL protein activity can be measured for changes in character by an appropriate erythrocyte binding assay.

Of course, due to the degeneracy of the genetic code, one of ordinary skill in the art will immediately recognize that a large number of the nucleic acid molecules having a sequence at least 70%, 80%, 90%, 95%, or 99% identical to a nucleic acid sequence shown herein in the Sequence Listing will encode a polypeptide "having BAEBL polypeptide activity." In fact, since degenerate variants of these nucleotide sequences all encode the same polypeptide, this will be clear to the skilled artisan even without performing the above described comparison assay. It will be further recognized in the art that, for such nucleic acid molecules that are not degenerate variants, a reasonable number will also encode a polypeptide having BAEBL polypeptide activity. This is because the skilled artisan is fully aware of amino acid substitutions that are either less likely or not likely to significantly effect protein function (e.g., replacing one aliphatic amino acid with a second aliphatic amino acid).

For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in (Bowie, J.U. *et al.* 1990 *Science* **247**:1306-1310), wherein the authors indicate that proteins are surprisingly tolerant of amino acid substitutions.

### Polypeptides and Fragments

5           The invention further provides a BAEBL polypeptide having the amino acid sequence encoded by an open reading frame (ORF) of a wild-type BAEBL gene, or a peptide or polypeptide comprising a portion thereof (e.g., region II).

          It will be recognized in the art that some amino acid sequences of the BAEBL polypeptides can be varied without significant effect of the structure or function of the protein. If such differences in sequence are contemplated, it should be remembered that  
10           there will be critical areas on the protein which determine activity.

          Thus, the invention further includes variations of the BAEBL polypeptide which show substantial BAEBL polypeptide activity or which include regions of BAEBL protein such as the protein portions discussed below. Such mutants include deletions, insertions,  
15           inversions, repeats, and type substitutions. As indicated, guidance concerning which amino acid changes are likely to be phenotypically silent can be found in (Bowie, J.U. *et al.* 1990 *Science* **247**:1306-1310).

          Thus, the fragment, derivative or analog of the polypeptide of the invention may be (i) one in which one or more of the amino acid residues are substituted with a conserved or  
20           non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which additional amino acids are fused to the mature polypeptide, such as a fusion peptide or leader or secretory sequence or a sequence which is employed for purification of  
25           the mature polypeptide or a proprotein sequence. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

          As indicated, changes are preferably of a minor nature, such as conservative amino acid substitutions that do not significantly affect the folding or activity of the protein (see Table A).



**Table A** Conservative Amino Acid Substitutions

<b>Aromatic</b>	Phenylalanine Tryptophan Tyrosine
<b>Ionizable: Acidic</b>	Aspartic Acid Glutamic Acid
<b>Ionizable: Basic</b>	Arginine Histidine Lysine
<b>Nonionizable Polar</b>	Asparagine Glutamine Serine Threonine
<b>Nonpolar (Hydrophobic)</b>	Alanine Glycine Isoleucine Leucine Proline Valine
<b>Sulfur Containing</b>	Cysteine Methionine

Amino acids in the BAEBL polypeptides of the present invention that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, 1989 *Science* 244:1081-1085). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as changes in erythrocyte binding character.

The polypeptides of the present invention are conveniently provided in an isolated form. By "isolated polypeptide" is intended a polypeptide removed from its native environment. Thus, a polypeptide produced and/or contained within a recombinant host cell is considered isolated for purposes of the present invention.

Also intended as an "isolated polypeptide" are polypeptides that have been purified, partially or substantially, from a recombinant host cell or a native source. For example, a recombinantly produced version of the BAEBL polypeptide can be substantially purified by the one-step method described in Smith and Johnson, 1988 *Gene* 67:31-40.

The polypeptides of the present invention include a polypeptide comprising a polypeptide shown herein in the Sequence Listing; as well as polypeptides which are at

least 70% identical, and more preferably at least 80%, 90%, 95% or 99% identical to those described above and also include portions of such polypeptides.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a reference amino acid sequence of an BAEBL polypeptide is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of the BAEBL polypeptide. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequence shown herein in the Sequence Listing can be determined conventionally using known computer programs such as the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, Wis. 53711). When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference amino acid sequence and that gaps in homology of up to 5% of the total number of amino acid residues in the reference sequence are allowed.

#### **Pharmaceutical Formulations and Modes of Administration**

The compounds of this invention can be employed in admixture with conventional excipients, i.e., pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral, enteral (e.g., oral) or topical application which do not deleteriously react with the active compounds. Suitable pharmaceutically acceptable carriers include but are not limited to water, salt solutions, alcohols, gum arabic, vegetable oils, benzyl alcohols,

polyethylene glycols, gelatine, carbohydrates such as lactose, amylose or starch, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and diglycerides, pentaerythritol fatty acid esters, hydroxy methylcellulose, polyvinyl pyrrolidone, etc. The pharmaceutical preparations can be sterilized and if desired  
5 mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, flavoring and/or aromatic substances and the like which do not deleteriously react with the active compounds. They can also be combined where desired with other active agents, e.g., vitamins.

10 For parenteral application, which includes intramuscular, intradermal, subcutaneous, intranasal, intracapsular, intraspinal, intrasternal, and intravenous injection, particularly suitable are injectable, sterile solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions, or implants, including suppositories. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose  
15 containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

20 For enteral application, particularly suitable are tablets, dragees, liquids, drops, suppositories, or capsules. The pharmaceutical compositions may be prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants  
25 (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid  
30 preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous

vehicles (e.g., almond oil, oily esters, ethyl alcohol or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer salts, flavoring, coloring and sweetening agents as appropriate. A syrup, elixir, or the like can be used wherein a sweetened vehicle is employed.

5           Sustained or directed release compositions can be formulated, e.g., liposomes or those wherein the active compound is protected with differentially degradable coatings, e.g., by microencapsulation, multiple coatings, etc. It is also possible to freeze dry the new compounds and use the lyophilizates obtained, for example, for the preparation of products for injection.

10           For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be  
15           determined by providing a valve to deliver a metered amount. Capsules and cartridges of, e.g., gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

          For topical application, there are employed as non-sprayable forms, viscous to semi-solid or solid forms comprising a carrier compatible with topical application and having a  
20           dynamic viscosity preferably greater than water. Suitable formulations include but are not limited to solutions, suspensions, emulsions, creams, ointments, powders, liniments, salves, aerosols, etc., which are, if desired, sterilized or mixed with auxiliary agents, e.g., preservatives, stabilizers, wetting agents, buffers or salts for influencing osmotic pressure, etc. For topical application, also suitable are sprayable aerosol preparations wherein the  
25           active ingredient, preferably in combination with a solid or liquid inert carrier material, is packaged in a squeeze bottle or in admixture with a pressurized volatile, normally gaseous propellant, e.g., a freon.

          The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may  
30           for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration.

### Reduced Binding to Gerbich Erythrocytes.

A new gene (*baebl*) of the DBL-EBP family of *Plasmodium* receptor proteins with erythrocyte specificity different from that of EBA-175 has been studied in *P. falciparum*. The exon/intron structure of BAEBL is similar to that of other DBL-EBL (Adams, J.H. *et al.* 1992 *PNAS USA* 89:7085-7089). The difference between two DBL-EBP from *P. falciparum* and *P. vivax/P. knowlesi* is the duplication of region II (Adams, J.H. *et al.* 1992 *PNAS USA* 89:7085-7089). Despite the similarity of BAEBL and EBA-175 in their requirement for sialic acid on erythrocyte proteins, the specificity of BAEBL for receptors on the erythrocyte surface differs from that of EBA-175. The differences are two-fold. First, BAEBL, but not EBA-175, binds to En(a-) erythrocytes that lack glycophorin A. Second, Gerbich erythrocytes (that have an altered glycophorin C and absent glycophorin D) bind BAEBL much more weakly than normal erythrocytes, but bind EBA-175 normally. Thus, the specificity of these two parasite receptor molecules differs, suggesting alternative pathways for invasion. These data indicate at least two different sialic acid-dependent pathways for invasion.

In preliminary experiments, BAEBL was not detected in the parasite supernatant absorbed with glycophorin C/D null erythrocytes of the Leach phenotype. BAEBL, however, was never detected in the eluate of these erythrocytes. In addition, 2 and 8  $\mu$ l of Leach erythrocytes, but not normal erythrocytes, removed BAEBL from parasite supernatant. The failure to elute BAEBL and its reduction with small numbers of Leach erythrocytes suggest proteolysis.

The Gerbich phenotype is found at high allele frequencies (50%) in some regions of Papua New Guinea (Booth, P.B. *et al.* 1982 *Hum Hered* 32:385-403). The mutation consists of a deletion of exon 3 in glycophorin C that leads to truncated glycophorin C and absent glycophorin D (Serjeantson, S.W. *et al.* 1994 *Immunol Cell Biol* 72:23-27). Of interest to our study is the fact that these areas co-localize to hyperendemic areas of malaria. Previously, Serjeantson (Serjeantson, S. W. 1989 *Papua New Guinea Med J* 32:5-9) found reduced frequency of heavy infections with *P. falciparum* and *P. vivax* with the Gerbich phenotype. The common mutation in erythrocyte band 3, resulting in ovalocytosis, was not described at the time of the study by Serjeantson (Serjeantson, S. W. 1989 *Papua New Guinea Med J* 32:5-9) and may have influenced the results. Pasvol *et al.* (Pasvol, G. *et al.* 1984 *Lancet* April 21, 907-908) found reduced invasion of Gerbich erythrocytes, but

these were also ovalocytic (Aanstee, D.J. *et al.* 1984 *Biochem J* 218:615-619). It is possible that hereditary ovalocytosis caused by band 3 mutations was influencing the invasion and frequency of infection. It is critical to restudy these groups now that the two mutations can be separated by molecular techniques. We were unable to find any reduction in invasion of Gerbich erythrocytes by the two *P. falciparum* clones, Dd2 and Dd2/Nm, but other parasite clones could be affected. It is known that Dd2 switched to the sialic acid-independent pathway (Dd2/Nm) after modification of the *eba-175* gene (Dolan, S.A. *et al.* 1990 *J Clin Invest* 86:618-624), suggesting that Dd2 lacking EBA-175 could not invade using BAEBL. Alternatively, the effect on invasion may be too subtle to be detected by the invasion assay as performed.

The tantalizing possibility remains that the high frequency of the Gerbich phenotype is selected for as a result of reduced invasion by *P. falciparum* mediated by the BAEBL receptor. The Gerbich mutation, like Duffy negativity in West Africa, appears to be going to fixation in these communities. The difference between the Gerbich mutation and Duffy negativity is that the Gerbich negative erythrocytes are still infected by *P. falciparum* while Duffy negative erythrocytes are refractory to *P. vivax*. Still, the parasite receptors for these two blood groups are envisioned as being immunogens to prevent malaria because mutation in the host molecules leads to no infection (*P. vivax* and Duffy negativity) or reduced infection (*P. falciparum* and Gerbich mutation).

#### Structure of the *baebl* Gene.

BAEBL is predicted to have two cysteine-rich domains (regions II and VI), a transmembrane region, and a cytoplasmic region. This structure is characteristic of all DBL-EBP genes (Adams, J. H. *et al.* 1992 *PNAS USA* 89:7085-7089). The sequence was obtained from the Sanger Centre chromosome 13 genomic sequence of *P. falciparum* clone 3D7 ([http://www.sanger.ac.uk/Projects/P\\_falciparum/](http://www.sanger.ac.uk/Projects/P_falciparum/)). The sequence of the gene was also determined from genomic and cDNA sequences of Dd2/Nm. The exon/intron structure for Dd2/Nm was identical to that of EBA-175 in that it had four exons: one for the extracellular domain, one for the transmembrane domain, and two encoding the cytoplasmic region (Fig. 1). The extracellular exon 1 of Dd2/Nm was identical to the genomic sequence of 3D7 from the Sanger Centre except for three changes in region II (I185V, N239S, and K261T; Dd2/Nm amino acid number from GenBank: AF332918/3D7 Sanger Centre chromosome 13). Using microsatellites and a genetic cross between Dd2 and HB3 (Su, X. *et al.* 1999

*Science* 286:1351-1353), *baebl* was localized to the end of chromosome 13 close to marker C13M51.

#### **Localization and Expression of BAEBL.**

Antibodies to the two cysteine-rich domains (regions II and VI) for BAEBL of Dd2/Nm were used to determine localization and expression of the protein. Antibodies to regions II and VI localize to the same organelle as EBA-175 (Fig. 2A, B), which was previously shown to localize to the micronemes (Sim, B.K.L. *et al.* 1992 *Mol Biochem Parasitol* 51:157-160). Furthermore, immunolocalization of RAP-1, a protein found in rhoptries, another apical organelle of merozoites, shows that BAEBL is adjacent to but not overlapping RAP-1 (Fig. 2C, D). This distribution is consistent with the localization of BAEBL within micronemes, a distribution identical to EBA-175 and the *P. knowlesi* Duffy binding protein (Adams, J.H. *et al.* 1990 *Cell* 63:142-153; Sim, B.K.L. *et al.* 1992 *Mol Biochem Parasitol* 51:157-160). The fact that antisera against two different regions of BAEBL showed identical localization within the parasite indicate that the antisera are not cross-reacting with another protein.

To study the molecular characteristics of BAEBL, we used methods developed for the production of soluble, metabolically labeled erythrocyte binding proteins (see Examples). Antisera to regions II and VI immunoprecipitated a protein of approximately 148 kDa; antibodies to region II also immunoprecipitated two lower molecular weight proteins (129 kDa and 117 kDa). The proof that the two 135-kDa proteins were identical derived from studies of immunoabsorption with one sera followed by immunoprecipitation with the second sera (Fig. 3). The same 135-kDa protein was removed by both sera, indicating that the antisera to the two regions of BAEBL were not cross-reacting with another protein. The two lower molecular weight proteins identified by anti-region II, but not by anti-region VI, resulted from immunoprecipitation of proteolytic products of BAEBL that contained region II but not region VI.

#### **Erythrocyte Binding Specificity.**

We have developed a new assay for measuring binding of BAEBL to erythrocytes. Previously, EBA-175 was identified in parasite proteins bound and eluted from human erythrocytes. Its identification was based on the fact that it was the most abundant and the highest molecular weight protein eluted from these erythrocytes. Lower molecular weight

proteins may be proteolytic fragments of EBA-175 or products of different genes. To positively identify BAEBL, we immunoprecipitated BAEBL from proteins eluted from erythrocytes with anti-region II and anti-region VI.

5 We also modified the protocol in that we identified proteins removed from the supernatant by erythrocyte absorption. We determined the different quantities of erythrocytes required to remove BAEBL from the parasite supernatant. It was found that 25  $\mu$ l of packed erythrocytes slightly reduced the quantity of immunoprecipitated BAEBL from 50  $\mu$ l of parasite supernatant. The protein was largely removed by absorbing twice with 50  $\mu$ l of packed erythrocytes for some culture supernatants and by only one absorption with 50  $\mu$ l for other supernatants. Therefore, on some samples, we absorbed with 25  $\mu$ l, 50  $\mu$ l once, and 50  $\mu$ l twice for comparison between normal erythrocytes and mutant erythrocytes or enzyme-treated erythrocytes. The protein was also eluted from the first 50  $\mu$ l of packed erythrocytes used for absorption. This set the conditions for absorbing and eluting BAEBL and demonstrated that BAEBL may be a parasite receptor for binding erythrocytes.

15 To determine the specificity of binding, we studied binding to neuraminidase-and trypsin-treated human erythrocytes and human erythrocytes with various genetically modified blood groups. Both enzymes eliminated the binding of BAEBL to human erythrocytes (Fig. 4), indicating that the erythrocyte receptor required sialic acid attached to a peptide backbone and must therefore be a sialoglycoprotein. This failure of neuraminidase-and trypsin-treated erythrocytes to bind was identical to EBA-175 (Fig. 4). To determine whether BAEBL was binding to the carbohydrates on the erythrocyte receptor, we performed competitive inhibition with Neu5Ac( $\alpha$ 2-3) lactosialic and Neu5Ac( $\alpha$ 2-6)lactosialic acid at 1  $\mu$ M, 10  $\mu$ M, 100  $\mu$ M, and 1000  $\mu$ M. We determined that 25 neither Neu5Ac( $\alpha$ 2-3) nor Neu5Ac( $\alpha$ 2-6) lactosialic acid inhibited the binding of BAEBL to human erythrocytes. These results indicate that BAEBL is binding either to a more complex polysaccharide or to a combination of sialic acid and a peptide backbone of an erythrocyte sialoglycoprotein.

30 To further define the binding specificity of BAEBL, we studied En(a-) erythrocytes, which lack glycophorin A. We found that EBA-175 failed to bind En(a-) erythrocytes as previously described (Sim, B.K.L. *et al.* 1994 *Science* 264:1941-1944). BAEBL, however, bound to these erythrocytes in a similar manner as to normal erythrocytes (Fig. 5). This



demonstrated that the binding specificity of BAEBL differed from that of EBA-175. S-s-U- erythrocytes that lacked glycophorin B bound both EBA-175 and BAEBL. Thus, neither glycophorin A nor B is the sole receptor for BAEBL.

#### **Abnormal Binding to Glycophorin C/D Mutant Erythrocytes.**

5           Another characterized sialoglycoprotein on the surface of human erythrocytes is glycophorin C/D (Reid, M.E. & Spring, F.A. 1994 *Transfusion Med* 4:139-146; Colin, Y. & Le Van Kim, C. 1995 in: *Blood Cell Biochemistry*, eds. Cartron, J. P. & Rouger, P. Plenum Press, New York pp. 331-350). Its peptide backbone is completely unrelated to glycophorins A and B but, like these, it has a mucin-like region of serines and threonines  
10           for O-linked sugars at the N-terminus of the protein. Both glycophorin C and glycophorin D are encoded by the same gene with use of alternative start codons. Glycophorin C, the full-length protein, contains one N-linked glycan. There are three mutations of the *glycophorin C/D* gene that lack high-incidence antigens (Colin, Y. & Le Van Kim, C. 1995 in: *Blood Cell Biochemistry*, eds Cartron, J.P. & Rouger, P. Plenum Press, New York pp.  
15           331-350). Leach erythrocytes are null for these proteins; Gerbich and Yus erythrocytes contain exon 3 and 2 deletions, respectively, that lead to a shortened glycophorin C and absent glycophorin D. Both Gerbich and Yus cells have abnormal N-linked glycosylation of the truncated form of glycophorin C (Reid, M.E. & Spring, F.A. 1994 *Transfusion Med* 4:139-146).

20           We screened for the binding of BAEBL to erythrocytes of the Gerbich (-2, -3, -4) and Yus (-2, -3, -4) phenotype that had been frozen as pellets in liquid nitrogen. BAEBL had reduced binding to Gerbich and Yus erythrocytes. These differences were consistent for pellet-frozen erythrocytes from different donors. EBA-175 bound normally to these erythrocytes.

25           Because the quality of the pellet-frozen erythrocytes was unpredictable, we obtained fresh blood from a person with the Gerbich mutation. In two separate experiments, we found that it required twice as many Gerbich cells to remove BAEBL from the culture supernatant compared with normal erythrocytes (Fig. 6A). In contrast to BAEBL, EBA-175 bound equally well to Gerbich and normal erythrocytes (Fig. 6C). This difference  
30           between normal and Gerbich erythrocytes for absorption of BAEBL was similar to the results obtained with pellet-frozen erythrocytes as described above.

BAEBL was eluted from normal but not Gerbich erythrocytes, indicative of its poor binding to Gerbich erythrocytes (Fig. 6B). BAEBL also did not elute from neuraminidase-treated normal erythrocytes. These results are indicative of poor binding of BAEBL to Gerbich erythrocytes. In contrast to BAEBL, EBA-175 was eluted from both Gerbich and normal erythrocytes (Fig. 6D).

#### Invasion of Gerbich Erythrocytes.

*P. falciparum* clones Dd2 and Dd2/Nm invaded Gerbich erythrocytes at the same rate as normal erythrocytes (Table 1). Dd2 but not Dd2/Nm had markedly reduced invasion into neuraminidase-treated erythrocytes as described previously (Dolan, S.A. *et al.* 1990 *J Clin Invest* 86:618-624).

**Table 1.** Invasion Rate of *P. falciparum* Into Gerbich Erythrocytes

Red-cell type	<i>P. falciparum</i> clones	
	Dd2, %	Dd2/Nm, %
Normal	3.7*	1.6
Gerbich	3.0	1.8
Neuraminidase-treated normal	0	1.8
Rhesus	0	0

\* Percentage of ring-infected erythrocytes.

#### EXAMPLE 1

##### Structure of BAEBL.

The sequence of BAEBL was identified (Adams J.H., *et al.* 2001 *Trends Parasitol* 17:17297-17299) from cDNA (GenBank No. N97830) deposited by D. Chakrabarti and from the database supplied by Sanger for chromosome 13 (>MAL13\_001500, December 27, 2000). Based on this sequence, we sequenced the *P. falciparum* clone, Dd2/Nm (Dolan, S.A. *et al.* 1990 *J Clin Invest* 86:618-624) from genomic DNA (GenBank No. AF332918). The exon/intron boundaries were defined by RT-PCR of the *P. falciparum* clone Dd2/Nm (GenBank No. AF332919). Primers use for Dd2/Nm sequencing were:

f1, 5'-AGACCAATAAATTATATATAATGAAAGGA-3' (SEQ ID NO: 3)

and 5'-TTTAAACTTTTCCATTGTTTCTAAACG-3' (SEQ ID NO: 4);

f2, 5'-ATAAATTTAATTCACCTTCCGAAAATGA-3' (SEQ ID NO: 5)

and 5'-AAAACAATCTCTTCTTTCCATCAAG-3' (SEQ ID NO: 6);

f3, 5'-TTTATAGGTGATGATATGGATTTTGG-3' (SEQ ID NO: 7)

and 5'-TCGTAAATGTTCCAGTACAATTCCT-3' (SEQ ID NO: 8);  
 f4, 5'-CAAATGGAGGTTTTGACGAAGTTG-3' (SEQ ID NO: 9)  
 and 5'-TAAGTACTGCTGACATTACTTTCCA-3' (SEQ ID NO: 10);  
 f5, 5'-GGAGCTTCAATATATGAGGCGCA-3' (SEQ ID NO: 11)  
 5 and 5'-ATATCTTCATATTCATTTGGACTCTC-3' (SEQ ID NO: 12);  
 f6, 5'-TGAGTCATTTAAGGTAGAATGTAAGA-3' (SEQ ID NO: 13)  
 and 5'-GGAAGTTTCCGAATGTCCATTCGT-3' (SEQ ID NO: 14);  
 f7, 5'-TAAATGAACAACAAAGTGGAAGGA-3' (SEQ ID NO: 15)  
 and 5'-ATTCTCAATTTGCGTTATATATTGATG-3' (SEQ ID NO: 16);  
 10 f8, 5'-AGTTCCTTCAGAGGATAATACCCA-3' (SEQ ID NO: 17)  
 and 5'-CTTGATTGACCCTCGCTTTTAAAC-3' (SEQ ID NO: 18);  
 f9, 5'-ACTAAAAGAGTAAGGGAGGAAATAAT-3' (SEQ ID NO: 19)  
 and 5'-TATAAAATACATTGAATTATTTAACTATTG-3' (SEQ ID NO: 20). PCR  
 from total RNA untreated with reverse transcriptase never produced PCR-amplified  
 15 products. Oligonucleotides 5'-ATTCCTTATTTTGCTGCTGGAGGT-3' (SEQ ID NO: 21)  
 and 5'-AAGTTGCTTCTATATTAGATTCTCCT-3' (SEQ ID NO: 22) were also used to  
 sequence fragment f9. Only the 3'-region was sequenced for cDNA to determine the  
 precise location of the intron/exon boundaries.

#### Antisera.

20 Antisera to BAEBL region II and region VI of Dd2/Nm were generated by  
 immunization of rats with a DNA vaccine using the vector VR1050 (kindly supplied by  
 Stephen Hoffman, Naval Medical Research Center, Silver Spring, MD) that contains the  
 T cell epitopes P2P30 from tetanus toxoid. Region II and region VI gene fragments of  
 BAEBL were amplified from *P. falciparum* clone Dd2/Nm and cloned into VR1050 vector,  
 25 previously described as VR1012tPAP2p30 by Becker *et al.* (Becker, S.I. *et al.* 1998 *Infect*  
*Immun* 66:3457-3461) but now renamed VR1050. The inserts for regions II and VI of  
 Dd2/Nm spanned from amino acids Q141 to I756 and K1046 to S1132, respectively  
 (GenBank No. AF332918). Rats were immunized intradermally with 500 µg of DNA at 3-  
 week intervals for a total of four immunizations. Sera were obtained from the rats a week  
 30 after the fourth immunization.

Rabbit anti-region II of EBA-175 (KLS14) was a kind gift of David Narum and  
 Kim Lee Sim (EntreMed, Rockville, MD). Mouse anti-RAP-1 monoclonal antibody

7H8/50 [MRA-79, Malaria Research and Reference Reagent Resource (MR4) Center] was a kind gift of Allan Saul (Queensland Institute of Medical Research, Brisbane, Australia).

#### **Erythrocytes Used in the Studies.**

5 Blood was collected in 10% citrate-phosphate-dextrose (vol/vol) and stored for up to 4 weeks at 4°C. At the time of study, the erythrocytes were washed three times in incomplete media (RPMI-1640; Life Technologies, Rockville, MD) with 25 mM HEPES and 0.36 mM hypoxanthine (Sigma, St. Louis, MO). For neuraminidase treatment, 5.5 ml of a 5% (vol/vol) suspension of the washed human erythrocytes in incomplete media were incubated twice with 3 milliunits of neuraminidase (*Vibrio cholerae*; CalBiochem, La Jolla, CA) for 2 hr at 37°C each time. For trypsin treatment, washed human erythrocytes were incubated with 1 mg/ml of tosyl-phenylalanine-chloromethyl-ketone-treated trypsin (Sigma) for 2 hr at 37°C. After trypsin treatment, the cells were washed once in incomplete medium and incubated with 2 mg/ml soybean trypsin inhibitor (Sigma) for 10 min at room temperature. The cells were washed twice before use in a study.

15 The glycophorin A and glycophorin B null erythrocytes [En(a-) and S-s-U-, respectively] and the glycophorin D null/glycophorin C modified erythrocytes (Gerbich cells) were frozen within a few days of receipt and thawed by the Red Cross method (Mallory, D. ed. 1993 *Immunohematology Methods and Procedures* American Red Cross, National Reference Laboratory, Rockville, MD, pp. 125-1-125-2). Blood from a Gerbich donor was collected in 10% (vol/vol) anticoagulant citrate-phosphate-dextrose.

20 Other glycophorin C/D mutant cells (Leach, Gerbich, and Yus cells) had been stored in liquid nitrogen as frozen pellets (Judd, W.J. 1994 in: *Methods in Immunohematology* Montgomery Scientific Publications Durham, NC, pp. 188-190) and thawed directly into PBS at 37°C.

#### **25 Metabolic Labeling of Parasite Proteins.**

Soluble, metabolically labeled parasite proteins were obtained from culture supernatant of schizont-infected erythrocytes that released merozoites in the absence of uninfected erythrocytes. The parasites were left to lyse and release proteins into the culture supernatant. The Dd2/Nm clone of *P. falciparum* was cultured as previously described (Kaneko, O. *et al.* 2000 *Mol Biochem Parasitol* 110:135-14) with the following exceptions. Schizont-infected erythrocytes ( $5 \times 10^7$  per ml of culture medium) were used during the

metabolic labeling. The culture supernatant was ultracentrifuged in a Beckman Optima TLX Ultracentrifuge (Beckman, Fullerton, CA) at 40,000 rpm (98,600×g) for 10 min at 4°C before storage at -70°C.

#### **Immunoprecipitation.**

5 Proteins in the supernatant and in the diluted eluate were immunoprecipitated as previously described (Kaneko, O. *et al.* 2000 *Mol Biochem Parasitol* 110:135-146) with the following exceptions. The supernatant (50 µl) was diluted into 250 µl of NETT (50 mM Tris, pH 7.4, 150 mM NaCl, 1 mM EDTA, and 0.5% Triton X-100) supplemented with 0.5% bovine serum albumin (BSA; ICN, Aurora, OH). To determine whether the proteins  
10 immunoprecipitated by anti-BAEBL region II and anti-BAEBL region VI are identical, we preabsorbed with one antisera and immunoprecipitated with the other. Radiolabeled supernatant (50 µl) was preabsorbed with protein A-Sepharose as previously described (Kaneko, O. *et al.* 2000 *Mol Biochem Parasitol* 110:135-146). The supernatant was incubated with 10 µl of anti-BAEBL region II or 10 µl of anti-BAEBL region VI for 2 hr at  
15 4°C. Protein G-Sepharose (40 µl; 50% vol/vol) was added to remove the immune complexes. Supernatant was split into two equal volumes and immunoprecipitated with 5 µl of anti-BAEBL region II and 5 µl anti-BAEBL region VI as described above.

#### **Modified Erythrocyte Binding Assay.**

Erythrocyte binding assays were developed for metabolically labeled proteins (as  
20 described above) that bind erythrocytes. The original assay required that parasite proteins be bound and eluted from some erythrocytes and not from others. In the original study (Camus, D. & Hadley, T.H. 1985 *Science* 230:553-556), the major protein eluted from the erythrocytes was EBA-175. Lower molecular proteins could be proteolytic fragments of EBA-175 or other proteins. Furthermore, this assay is insensitive for less abundant  
25 proteins. Therefore, we have developed a new assay that depends on the identification of BAEBL with two antisera against different regions of BAEBL and its removal from the culture supernatant by human erythrocytes. The parasite protein can also be identified and quantified by elution of bound protein from erythrocytes followed by immunoprecipitation. It is then possible to study its specificity for erythrocyte receptors with erythrocytes lacking  
30 various proteins or with enzymatically modified erythrocytes. First, we determined the quantity of erythrocytes that would remove the majority of BAEBL and used this quantity

with erythrocytes of various types (enzyme-modified erythrocytes and erythrocytes genetically deficient in membrane proteins) to determine the erythrocyte specificity of BAEBL. We found that one or two absorptions with a volume of packed erythrocytes equal to the volume of metabolically labeled supernatant were required to remove BAEBL from the supernatant, depending on the concentration of BAEBL in the supernatant.

Elution from erythrocytes of bound parasite proteins was performed as described previously (Kaneko, O. *et al.* 2000 *Mol Biochem Parasitol* 110:135-146). Parasite proteins were eluted only from the erythrocytes of the first adsorption. The parasite proteins were eluted as previously described. Because of the adverse effect of high salt on immunoprecipitation, the eluate was diluted 5 fold (vol/vol) in NETT with 0.5% BSA prior to immunoprecipitation.

#### **Competitive Inhibition Assay.**

An inhibition assay was conducted in the presence of Neu5Ac( $\alpha$ 2-3) lactosialic acid or Neu5Ac( $\alpha$ 2-6) lactosialic acid (Sigma). Metabolically labeled parasite supernatant (50  $\mu$ l) was preincubated with 1, 10, 100, or 1000  $\mu$ M in 15  $\mu$ l of the aforementioned carbohydrates for 1 hr at room temperature. Packed erythrocytes (50  $\mu$ l) were added to the mixture. The erythrocyte binding assay was conducted as described above.

#### **Immunolocalization of BAEBL.**

The methods for immunolocalization of BAEBL by confocal microscopy were performed as previously described (Kaneko, O. *et al.* 2000 *Mol Biochem Parasitol* 110:135-146) with the following modifications. The blocking buffer consisted of PBS (pH 7.4) containing 0.1% Triton X-100 (Bio-Rad, Hercules, CA) and 2.5 mg/ml normal goat serum (Jackson ImmunoResearch Laboratories, West Grove, PA). The secondary antisera consisted of Alexa 488-conjugated goat anti-rat IgG and Alexa 594-conjugated goat anti-rabbit IgG (Molecular Probes, Eugene, OR) diluted 1:500 in blocking buffer. For anti-quenching, we mounted labeled parasites in Prolong Antifade (Molecular Probes).

#### **Invasion Assay.**

Pre-washed A<sup>+</sup> human erythrocytes treated with neuraminidase and A<sup>+</sup> human erythrocytes of the Gerbich type (-2, -3, -4) were tested for invasion by *P. falciparum* clones Dd2 and Dd2/Nm (Dolan, S.A., *et al.* 1990 *J Clin Invest* 86:618-624) as described previously (Kaneko, O. *et al.* 1999 *Exp Parasitol* 93:116-119). Rhesus (*Macaca mulatta*)

erythrocytes that are resistant to invasion by *P. falciparum* were used as a control for normal erythrocytes introduced with the parasitized erythrocytes.

## **EXAMPLE 2**

*Plasmodium falciparum* has evolved great flexibility in its invasion pathways, in part as a result of multiple copies of the Duffy binding-like (DBL) family of erythrocyte-binding ligands, three of which have different red blood cell (RBC) receptor specificities (Sim, B.K.L. *et al.* 1994 *Science* 264:1941-1944; Mayer D.C.G. *et al.* 2001 *PNAS USA* 98:5222-5227). This is in contrast to *P. vivax*, which has a single copy of the DBL family member that recognizes the Duffy blood group antigen, explaining the resistance to *P. vivax* infection by humans who lack the Duffy blood group (Chitnis C.E. *et al.* 1994 *J Exp Med* 180:497-506). The multiplicity of *P. falciparum* invasion pathways explains the lack of RBC refractory to invasion. We now describe another mechanism for recognition of different molecules on the RBC by *P. falciparum*, namely, amino acid polymorphisms in the DBL gene, BAEBL, that lead to different RBC specificities (Adams J.H., *et al.* 2001 *Trends Parasitol* 17:17297-17299).

Initial characterization of RBC receptors recognized by BAEBL from three different *P. falciparum* clones in three different laboratories suggested that each had a different RBC receptor (Sim, B.K.L. *et al.* 1994 *Science* 264:1941-1944; Thompson, J.K. *et al.* 2001 *Mol Microbiol* 41:47-58; Narum, D.L. *et al.* 2002 *Mol Biochem Parasitol* 119:159-168). We sequenced the *baebl* gene from eight parasite clones and found polymorphisms restricted to regions I and II of the molecules. In total, we sequenced region II of BAEBL from 11 clones from Papua New Guinea (PNG) and 13 clones from other parts of the world where *P. falciparum* malaria is highly prevalent. We observed five different sequence variants in region II with polymorphisms in four amino acid positions (Table 2). Unlike Africa where *P. falciparum* clones were introduced from Asia, PNG *P. falciparum* populations are isolated. Surprisingly, the same sequence variants occurred in PNG as in the rest of the world, suggesting that mutations leading to these polymorphisms occurred multiple times. Region II of *P. falciparum* DBL genes are duplicated forming the F1 and F2 domains (Adams J.H. *et al.* 1992 *PNAS USA* 89:7085-7089). All base substitutions in the erythrocyte binding domain of BAEBL occurred in the F1 domain, whereas mutations in EBA-175, another DBL gene of *P. falciparum*, were scattered throughout both F1 and F2

domains and appear not to alter the RBC binding specificity (Liang, H. & Sim, B.K. 1997 *Mol Biochem Parasitol* 84:241-245).

**Table 2** Position of Polymorphisms in BAEBL From Different Malaria-Endemic Regions

Clones	Origin	Region I *		Region II *			
		26	112	185	239	261	285
PNG2	PNG †	I (ATT)	L(CTT)	V(GTT)	S(AGT)	T(ACG)	K(AAA)
PNG3	PNG	I (ATT)	F(TTT)	I(ATT)	N(AAT)	R(AGG)	E(GAA)
PNG4	PNG	I (ATT)	L(CTT)	V(GTT)	S(AGT)	K(AAG)	K(AAA)
E12	PNG	I (ATT)	F(TTT)	I(ATT)	S(AGT)	K(AAG)	K(AAA)
1917	PNG	I (ATT)	L(CTT)	V(GTT)	S(AGT)	K(AAG)	K(AAA)
PNG13	PNG	I (ATT)	F(TTT)	V(GTT)	S(AGT)	K(AAG)	K(AAA)
PNG5	PNG	I (ATT)	L(CTT)	I(ATT)	S(AGT)	K(AAG)	K(AAA)
1905	PNG	I (ATT)	L(CTT)	V(GTT)	S(AGT)	K(AAG)	K(AAA)
PNG9-3	PNG	I (ATT)	L(CTT)	V(GTT)	S(AGT)	K(AAG)	K(AAA)
PNG9-1	PNG	I (ATT)	L(CTT)	V(GTT)	S(AGT)	T(ACG)	K(AAA)
PNG10-1	PNG	I (ATT)	L(CTT)	I(ATT)	S(AGT)	K(AAG)	K(AAA)
M24	Kenya	I (ATT)	F(TTT)	I(ATT)	N(AAT)	K(AAG)	K(AAA)
3D7	Africa?	I (ATT)	F(TTT)	I(ATT)	N(AAT)	K(AAG)	K(AAA)
Sc/d6	Sierra Leone	I (ATT)	L(CTT)	V(GTT)	S(AGT)	K(AAG)	K(AAA)
Fab9	Kwazulu	I (ATT)	L(CTT)	I(ATT)	N(AAT)	R(AGG)	E(GAA)
Dd2	Vietnam	I (ATA)	L(CTT)	V(GTT)	S(AGT)	T(ACG)	K(AAA)
Camp	Malaysia	I (ATT)	L(CTT)	I(ATT)	N(AAT)	R(AGG)	E(GAA)
Dd2/Nm	Vietnam	I (ATT)	L(CTT)	V(GTT)	S(AGT)	T(ACG)	K(AAA)
T2/c6	Thailand	I (ATT)	F(TTT)	I(ATT)	N(AAT)	K(AAG)	K(AAA)
MT/S-1	Asia	I (ATT)	F(TTT)	I(ATT)	S(AGT)	K(AAG)	K(AAA)
HB3	Honduras	I (ATT)	L(CTT)	V(GTT)	S(AGT)	K(AAG)	K(AAA)
PC49	S. America	I (ATT)	F(TTT)	I(ATT)	N(AAT)	K(AAG)	K(AAA)
DIV30	Brazil	I (ATT)	L(CTT)	V(GTT)	S(AGT)	T(ACG)	K(AAA)
PC26	S. America	I (ATT)	L(CTT)	V(GTT)	S(AGT)	T(ACG)	K(AAA)

\* For Regions I and II, numbers refer to the amino acids in the sequence of BAEBL from GenBank AF332918. Mutated bases and amino acids are shown in **bold**.

† PNG, Papua New Guinea.



We investigated the functional significance of polymorphisms in the erythrocyte-binding domain of BAEBL. We expressed region II of four polymorphic groups transiently on the surface of COS cells with the T8 vector (Buffet P.A. *et al.* 1999 *PNAS USA* 96:12743-12748). This was followed by an erythrocyte-binding assay as described previously (Chitnis C.E. *et al.* 1994 *J Exp Med* 180:497-506). Binding was performed with normal and enzyme-treated (trypsin and neuraminidase) erythrocytes and Gerbich-negative erythrocytes (exon 3 deletion of glycophorin C/D) (Sim, B.K.L. *et al.* 1994 *Science* 264:1941-1944; Mayer D.C.G. *et al.* 2001 *PNAS USA* 98:5222-5227). Each of the polymorphisms led to a different binding specificity as demonstrated by different binding patterns to enzyme-treated and Gerbich-negative RBC (Table 3). Furthermore, a single base change led to a change in amino acid and RBC specificity (e.g., VSTK to VSKK). Such polymorphism in sequence and receptors was described for influenza hemagglutinin, where a single base mutation changed the amino acid and the specificity of binding to sialic acid (Rogers, G.N. *et al.* 1983 *Nature* 304:76-78).

**Table 3** Binding Patterns of BAEBL Variants to Enzyme-Treated and Gerbich-Negative RBC

Region II variants*	Normal RBC			Gerbich-negative RBC ‡
	Untreated †	Trypsin ‡	Neuraminidase ‡	
VSTK	65	0	0	0.9%
VSKK	58	0	90%	112%
ISKK	59	92%	0	96%
INRE	67	114%	110%	100%

\* Region II of BAEBL expressed in COS cells is from amino acid 143 to 606 (GenBank AF332918) and contains the mutations shown at the positions delineated in Table 2.

† COS cells with five or more attached RBC were counted and the total per coverslip recorded.

‡ Data from enzyme-treated and Gerbich-negative RBC are expressed as the percentage of binding to normal, untreated RBC.

The Gerbich-negative phenotype occurs at an allelic frequency of 50% in some regions of PNG (Booth, P.B. *et al.* 1982 *Human Hered* 32:385-403). It is tantalizing to postulate that the polymorphisms in BAEBL could be a coevolutionary adaptation to the

disappearance of the RBC receptor in Gerbich-negative individuals. Why then has the Gerbich-negative phenotype not been described in all the geographic areas where mutations in region II of BAEBL have occurred? One possibility is that the Gerbich-negative phenotype could indeed be more widespread than previously described. Alternatively, such polymorphisms in BAEBL receptor specificity may be advantageous to the parasite independently of the Gerbich negative phenotype.

\*\*\*\*\*

While the present invention has been described in some detail for purposes of clarity and understanding, one skilled in the art will appreciate that various changes in form and detail can be made without departing from the true scope of the invention. All figures, tables, and appendices, as well as patents, applications, and publications, referred to above, are hereby incorporated by reference.

WHAT IS CLAIMED IS:

1. A vaccine composition comprising a polypeptide and a pharmaceutically acceptable vehicle, wherein the polypeptide comprises an amino acid sequence that encodes a BAEBL polypeptide or portion thereof.
- 5 2. A vaccine composition of Claim 1, wherein the polypeptide portion is an amino acid sequence that encodes a BAEBL region II or portion thereof.
3. A vaccine composition of Claim 2, wherein the polypeptide portion is selected from the group consisting of an amino acid sequence having the following number of consecutive amino acids taken from said BAEBL polypeptide:  
10 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120,  
15 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210,  
20 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300,  
25 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390,  
30 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444,

445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, and 584.

4. A vaccine composition of any of Claims 1-3 wherein said BAEBL polypeptide or portion thereof is defined as having the amino acid sequence of SEQ ID NO: 2 or portion thereof.

5. A vaccine composition of any of Claims 1-3 wherein said BAEBL polypeptide or portion thereof is defined as having at least 70%, 80%, 90%, 95%, or 99% identity to the amino acid sequence of SEQ ID NO: 2 or portion thereof.

6. A vaccine composition of any of Claims 1-3 wherein said BAEBL polypeptide or portion thereof is encoded by a polynucleotide defined as having at least 70%, 80%, 90%, 95%, or 99% identity to the open reading frame of SEQ ID NO: 1 or portion thereof.

7. A vaccine composition of any of Claims 1-3 wherein said BAEBL polypeptide or portion thereof is encoded by a polynucleotide which hybridizes at 42 degree C. in a solution comprising: 50% formamide, 5 times SSC (750 mM NaCl, 75 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 times Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1 times SSC at about 65 degree C, to a second polynucleotide having the polynucleotide sequence of SEQ ID NO: 1.

8. A vaccine composition of any of Claims 1-3, wherein said BAEBL polypeptide or portion thereof has a polymorphism selected from the group consisting of I at position 185, N at position 239, T at position 261, R at position 261, and E at position 285.

9. A vaccine composition of any of Claims 1-3 further comprising an adjuvant selected from the group consisting of QS-21, Detox-PC, MPL-SE, MoGM-CSF, TiterMax-

G, CRL-1005, GERBU, TERamide, PSC97B, Adjuvax, PG-026, GSK-1, GcMAF, B-alethine, MPC-026, Adjuvax, CpG ODN, Betafectin, Alum, and MF59.

10. A vaccine composition of any of Claims 1-3 further comprising a second polypeptide, wherein said second polypeptide comprises an amino acid sequence that encodes at least a portion of a Duffy binding protein or erythrocyte binding antigen-175 (EBA-175) of a malaria Plasmodium parasite.

11. A vaccine composition comprising a polynucleotide and a pharmaceutically acceptable vehicle, wherein the polynucleotide comprises a nucleic acid sequence that encodes a BAEBL polypeptide or portion thereof.

12. A vaccine composition of Claim 11, wherein the polypeptide portion is an amino acid sequence that encodes a BAEBL region II or portion thereof.

13. A vaccine composition of Claim 12, wherein the polypeptide portion is selected from the group consisting of an amino acid sequence having the following number of consecutive amino acids taken from said BAEBL polypeptide:

6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354,

355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372,  
373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390,  
391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408,  
409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426,  
5 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444,  
445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462,  
463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480,  
481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498,  
499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516,  
10 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534,  
535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552,  
553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570,  
571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, and 584.

14. A vaccine composition of any of Claims 11-13 wherein said BAEBL  
15 polypeptide or portion thereof is defined as having the amino acid sequence of SEQ ID NO:  
2 or portion thereof.

15. A vaccine composition of any of Claims 11-13 wherein said BAEBL  
polypeptide or portion thereof is defined as having at least 70%, 80%, 90%, 95%, or 99%  
identity to the amino acid sequence of SEQ ID NO: 2 or portion thereof.

20 16. A vaccine composition of any of Claims 11-13 wherein said BAEBL  
polypeptide or portion thereof is encoded by a polynucleotide which is identical to the open  
reading frame of SEQ ID NO: 1 or portion thereof.

17. A vaccine composition of any of Claims 11-13 wherein said BAEBL  
polypeptide or portion thereof is encoded by a polynucleotide defined as having at least  
25 70%, 80%, 90%, 95%, or 99% identity to the open reading frame of SEQ ID NO: 1 or  
portion thereof.

18. A vaccine composition of any of Claims 11-13 wherein said BAEBL  
polypeptide or portion thereof is encoded by a polynucleotide which hybridizes at 42  
degree C. in a solution comprising: 50% formamide, 5 times SSC (750 mM NaCl, 75 mM  
30 trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 times Denhardt's solution, 10%  
dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing

the filters in 0.1 times SSC at about 65 degree C, to a second polynucleotide having the polynucleotide sequence of SEQ ID NO: 1.

5           19.    A vaccine composition of any of Claims 11-13, wherein said BAEBL polypeptide or portion thereof has a polymorphism selected from the group consisting of I at position 185, N at position 239, T at position 261, R at position 261, and E at position 285.

          20.    A method of vaccinating a human against a malaria Plasmodium parasite comprising the step of administering the vaccine composition of any of the above claims to said human.

10           21.    The method of Claim 20 wherein said step of administration is by protein immunization.

          22.    The method of Claim 20 wherein said step of administration is by genetic immunization.

15           23.    Use of the vaccine composition of any of the above claims for the preparation of a medicament for vaccinating a human against a malaria Plasmodium parasite.

          24.    A method of vaccinating a human against a malaria Plasmodium parasite comprising the step of administering antibodies specific for the binding site of a BAEBL ligand in an amount sufficient to inhibit the ligand from binding red blood cells in the human.

20




FIG. 1A

FIG. 1B






FIG. 2A

FIG. 2B

FIG. 2C

FIG. 2D

3 / 8




FIG. 3





FIG. 4

4 / 8



**FIG. 5**






FIG. 7A
FIG. 7B
FIG. 7C

# FIG. 7A

SEQ ID NO: 1

## BAEBL NUCLEOTIDE SEQUENCE

# FIG. 7

tttataatat cgttttttta tgagcattat atcatataaa taattaatta aataaatata  
 tataatata taattataat tagaccaata aattatatat aatgaaggga tattttaata  
 tataattttt aattccttta atttttttat ataattgtaat aagaataaat gaatcaataa  
 taggtagaac actttataat agacaagatg aatcatcaga tatttcaagg gtaaatcac  
 ccgaattaaa taataatcat aaactaata tatatgattc agattacgaa gatgtaata  
 ataaattaat aaacagtttt gtagaaaata aaagtgtgaa aaaaaaagg tctttaagtt  
 ttataaataa taaaacaaaa tcatatgata taattccacc ttcataattca tataggaatg  
 ataaatttaa ttcactttcc gaaaatgaag ataatttctgg aaatacaaat agtaataatt  
 tcgcaaatac ttctgaaata tctattggaa aggataataa acaatatagc tttatacaga  
 aacgtactca ttgttttgct tgtggaataa aaagaaaatc aataaaatgg atatgtcgag  
 aaaacagtga gaaaattact gtatgtgttc ctgatagaaa aatacaacta tgtgttgcaa  
 atttttttaa ctcacgttta gaaacaatgg aaaagtttaa agaaatattt ttaatttctg  
 ttaatacaga agcaaaaatta ttataaaca aaatgaagg aaagatccc tcaatatatt  
 gtaatgaatt aagaaatagt ttttcagatt ttagaagttc atttataggt gatgatagg  
 attttgggtg taatacagat agagtcaaaag gatataataa tacgaagtcc tccgattatt  
 ataaggaaaa aaatgttgaa aaattaaata atatcaaaaa agaattggtgg gaaaaaata  
 aagcaaatat gtggaatcac atgatagtaa atcataaagg aaacataagt aaagaatgtg  
 ccataattcc cgcggaagaa cctcaaatat atctatggat aaaagaatgg aatgaaaaact  
 tcttgatgga aaagaagaga ttgtttttta atataaaga taagtgtgtt gaaaaaaaa

7/8

```

aatatgaagc atgttttggg ggatgtaggc ttccatgttc ttcatataca tcatttatga
aaaaaagtaa aacacaaaatg gaggttttga cgaacttgta taaaaagaaa aattcaggag
tgataaaaaa taatttttctg aatgatcttt ttaaaaaaaa taataaaaaa gatttagatg
attttttcaa aatgaaaaag gaatatgatg atttatgtga ttgcagatat actgctacta
ttattaaaag ttttctaaat ggtcctgcta aaatgatgt agatatgtga tcacaaatta
atgttaatga tcttcgaggg tttggatgta attataaaaag taataatgaa aaaagttgga
attgtactgg aacattttacg acaaaatttc ctggtacatg tgaaccccc agaagacaaa
ctttatgtct tggacgtaca tatcttttac atcgtggtca tgaggaagat tataaggaac
atttacttgg agcttcaata tatgagggcg aattattaaa atataaatat aaggaaaagg
atgaaaatgc attgtgtagt ataatacaaa atagttatgc agatttggca gatattatca
agggatcgga tataataaaa gattattatg gtaaaaaaat ggaagaaaat ttaataaaa
taacaaaaga taaaaaacgt aatgaagaat cttgaaagt tttcgtgaa aaatggtggg
atgaaaacaa ggagaatgta tggaaagtaa tgtcagcagt acttaaaaat aaggaaacgt
gtaaagatta tgataagttt caaaagattc ctcaattttt aagatggttt aaggaatggg
gagacgattt ttgtgagaaa agaaaagaga aaatatattc atttgagtca tttaaggtag
aatgtaagaa aaaagattgt gatgaaaata catgtaaaaa taaatgtagt gaataaaaa
aatggataga ttgaaaaaaa agtgaatatg agaaacaaagt tgataaatat acaaaagata
aaaataaaaa gatgtatgat aatattgatg aagtaaaaaa taaagaagcc aatgtttact
taaaagaaaa atccaaagaa tgtaaagatg taaatttcga tgataaaatt tttaatgaga
gtccaaatga atatgaagat atgtgtaaaa aatgtgatga aataaaatat ttaaatgaaa
ttaaatatcc taaaacaaaa cacgatatat atgatataga tacattttca gatacttttg
gtgatggaac gccaatagat attaatgcaa atataaatga acaacaaagt ggggaaggata
cctcaaatatc tggaaatagt gaaacatcag attcacgggt tagtcatgaa ccagaaagtg
atgctgcaat taatgtagaa aagttaagtgt gtgatgaaag ttcaagtgaa acaagaggaa
tattagatat taatgatcca agtgttacga acaatgtcaa tgaagttcat gatgcttcaa

```

FIG. 7B

8 / 8

atacacaagg tagtggtttca aatacttctg atataacgaa tggacattcg gaaagttccc  
 tgaatagaac aacgaatgca caagatatata aataggccg ttcaggaaat gaacaaagtg  
 ataatacaga aaatagttca cattctagt cattctcagg ataattcagg tctttgaca atcgacaag  
 ttccttcaga ggataatacc caaataacat atgattcaca aaacctcat agagatacac  
 ctaatgcatt agcatcttta ccatacagatg ataaaaattaa tgaatatagag ggtttcgatt  
 ctagtagaga tagtgaaaat ggtaggggtg atacaacatc aaatactcat gatgtacgtc  
 gtacgaatat agtaagtga agacgtgtga atagccatga tttattaga aacggaatgg  
 cgaataacaa tgcacatcat caatatataa cgcaaatga gaataatgga atcataagag  
 gacaagagga aagtgcgggg aatagtgtta attataaaga taatccaaag aggagtaatt  
 ttcctccga aaatgatcat aagaaaaata tacagggaata taattctaga gatactaaaa  
 gagtaaggga ggaataaatt aaattatcga agcaaaataa atgcaacaat gaataatcca  
 tggaatatgt tacctattct gacgaaaagga atagttcacc ggtccttgt tctagagaag  
 aaagaaaaga attatgttgt cagatttcag attattgttt aaaataattt aacttttatt  
 caattgaata ttataattgt ataaaatctg aaattaaaag tccagaatat aaatgtttta  
 aaagcgaggg tcaatcaagt atgtttcata tataatgaaa ataactaaat taaaataaat  
 atctaaaatt tccatttaaa tataatttaa ataatttatt tttttattat aaaatttatt  
 tgtttttttt tataaatgt ttattttttt cttacaggg attccttatt ttgctgctgg  
 aggtatttta gttgtaatag tcttactttt gagttcagca tctagaatgg ggaaaagggtt  
 acacgcataa gattaaaata atatgactat actttttaat ttttatatta ttatatatt  
 tattatgttt taacaagaat ataaattata ttatatattt atatgtatat atattttttt  
 atatagtaat gaagaatatg atataggaga atctaataa taagattaat tgaagaaaa  
 taattattta aataaaactat cgcgcatatg taagattaat ataaaaagt aaaatttcat  
 attaaataaaa taaatgaata tatatatata tatatatata tatatacatg  
 ttataataat tttaaatgtta tattatttac ttttttctca gttaatcaag aagtacaaga  
 gacaaacatt tcagattatt ccgagtacaa ttataatgaa aagaatatgt attaat

## SEQ ID NO:1

## BAEBL NUCLEOTIDE SEQUENCE

tttataatat cgttttttta tgagcattat atcatataaa taattaatta aataaatata  
 tatatatata taattataat tagaccaata aattatataat aatgaaagga tattttaata  
 tatatttttt aattccttta atttttttat ataattgtaat aagaataaat gaatcaataa  
 taggtagaac actttataat agacaagatg aatcatcaga tatttcaagg gtaaattcac  
 ccgaattaaa taataatcat aaaactaata tatatgattc agattacgaa gatgtaaata  
 ataaattaat aaacagtttt gtagaaaata aaagtgtgaa aaaaaaagg tctttaagtt  
 ttataaataa taaaacaaaa tcatatgata taattccacc ttcataattca tataggaatg  
 ataaatttaa ttcacttttc gaaaatgaag ataattctgg aaatacaaat agtaataatt  
 tcgcaaatat tcttgaaata tctattggaa aggataataa acaatatagc tttatacaga  
 aacgtactca tttgtttgct tgtggaataa aaagaaaatc aataaaatgg atatgtcga  
 aaaacagtga gaaaattact gtatgtgttc ctgatagaaa aatacaacta tgtgttgcaa  
 atttttttaa ctcacgttta gaaacaatgg aaaagtttaa agaatatttt ttaatttctg  
 ttaatacaga agcaaaatta ttatataaca aaaatgaagg aaaagatccc tcaatatttt  
 gtaatgaatt aagaaatagt ttttcagatt ttggaagttc atttataggt gatgatattg  
 attttgggtg taatacagat agagtcaaag gatataataa tacgaagttc tccgattatt  
 ataaggaataa aaatgttgaa aaattaaaata atatacaaaa agaattggtg gaaaaaaaata  
 aagcaaatgt gtggaatcac atgatagtaa atcataaagg aaacataagt aaagaatgtg  
 ccataattcc cgcggaagaa cctcaaatta atctatggat aaaagaatgg aatgaaaact  
 tcttgatgga aaagaagaga ttgtttttaa atataaaaga taagtgtgtt gaaaacaaaa  
 aatatgaagc atgttttggg ggatgtaggc ttccatgttc ttcataatata tcatatttga  
 aaaaaagtaa aacacaaatg gaggttttga cgaacttgta taaaaagaaa aattcaggag  
 tgataaaaaa taattttctg aatgatcttt ttaaaaaaaa taataaaaaat gatttagatg  
 attttttcaa aaatgaaaag gaatatgatg atttatgtga ttgcagatat actgctacta  
 ttattaaaag ttttctaatt ggtcctgcta aaaaatgatgt agatattgca tcacaaatta  
 atgttaatga tcttcgaggg tttggatgta attataaaag taataatgaa aaaagttgga  
 attgtactgg aacattttac aacaaatttc ctggtacatg tgaaccccc agaagacaaa  
 ctttatgtct tgacgtaca tatcttttac atcgtggtca tgagggaagat tataagggaac  
 atttacttgg agcttcaata tatgaggcgc aattattaaa atataaatat aaggaaaagg  
 atgaaaatgc attgtgtagt ataatacaaa atagttatgc agatttggca gatattatca  
 agggatcgga tataataaaa gattattatg gtaaaaaaat ggaagaaaat ttaaataaag  
 taaacaaaaga taaaaaacgt aatgaagaat ctttgaagat ttttcgtgaa aaatggtggg  
 atgaaaacaa ggagaatgta tggaaagtaa tgtcagcagt acttaaaaat aaggaaacgt  
 gtaaagatta tgataagttt caaaagattc ctcaattttt aagatggttt aagggaatggg  
 gagacgattt ttgtgagaaa agaaaagaga aaatatattc atttgagtca ttttaggtag  
 aatgtaagaa aaaagattgt gatgaaaata catgtaaaaa taaatgtagt gaataaaaa  
 aatggataga tttgaaaaaa agtgaatatg agaaacaagt tgataaatac acaaaagata  
 aaaataaaaa gatgtatgat aatattgatg aagtataaaa taaagaagcc aatgtttact  
 taaaagaaaa atccaaagaa tgtaaagatg taaatttoga tgataaaatt tttaatgaga  
 gtccaaatga atatgaagat atgtgtaaaa aatgtgatga aataaaatat ttaaatgaaa  
 ttaaatatcc taaaacaaaa cacgatatat atgatataga tacattttca gatacttttg  
 gtgatggaac gccaaataagt attaatgcaa atataaatga acaacaaagt gggaaggata  
 cctcaaatac tggaaatagt gaaacatcag attcaccggt tagtcatgaa ccagaaagtg  
 atgctgcaat taatgtagaa aagttaagtg gtgatgaaag ttcaagtga acaagaggaa  
 tattagatat taatgatcca agtgttacga acaatgtcaa tgaagttcat gatgcttcaa  
 atacacaagg tagtgtttca aatacttctg atataacgaa tggacattcg gaaagttccc  
 tgaatagaac aacgaatgca caagatatta aaataggccg ttcaggaaat gaacaaagtg  
 ataatacaga aaatagttca cattctagtg ataattcagg ttctttgaca atcggaacag  
 ttccttcaga ggataatacc caaaatacat atgattcaca aaacctcat agagatacac  
 ctaatgcatt agcatcttta ccatcagatg ataaaattaa tgaaatagag ggtttcgatt  
 ctagtagaga tagtgaaaat ggtaggggtg atacaacatc aaatactcat gatgtacgtc  
 gtacgaatat agtaagttag agacgtgtga atagccatga ttttattaga aacggaatgg  
 cgaataacaa tgcacatcat caatatataa cgcaaattga gaataatgga atcataagag  
 gacaagagga aagtgcgggg aatagtgtta attataaaga taatccaaag aggagtaatt  
 tttcctccga aaatgatcat aagaaaaata tacaggaata taattctaga gatactaaaa  
 gagtaaggga ggaataaatt aaattatcga agcaaaaata atgcaacaat gaattattcca  
 tggaattatt tacctattct cagcaaaagg atagttcacc gggctcttgt tctagagaag  
 aaagaaagaa attatgttgt cagatttcag attattgttt aaaaattttt aactttttat  
 caattgaata ttataattgt ataaaaatctg aaattaaaag tccagaatat aaatgtttta

Figure 7



```
aaagcgaggg tcaatcaagt atgtttcata tataatgaaa ataactaaat taaaataaat  
atctaaattt tccatttaaa tataatttaa ataatttatt tttttattat aaaatttatt  
tgtttttttt tatataatgt tttatttttt ctttacaggc attccttatt ttgctgctgg  
aggtatttta gttgtaatag tcttactttt gagttcagca tctagaatgg ggaaaagggt  
acacgcataa gattaaataa atatgactat actttttaat ttttatatta tttatattat  
tattatgttt taacaagaat ataaattata ttatatattt atatgtatat atattttttt  
atatagtaat gaagaatatg atataggaga atctaataa gaagcaactt ttgaagaaaa  
taattattta aataaaactat cgcgcataat taagattaat ataaaaagtg aaaatttcat  
attaaataaa taaatgaata tatatatata tatatatata tatatatata tatatacatg  
ttatataaat tttaatgtta tattatttac ttttttctca gttaatcaag aagtacaaga  
gacaaacatt tcagattatt ccgagtacaa ttataatgaa aagaatatgt attaat
```

Figure 7 cont'd

## SEQUENCE LISTING

<110> The Government of the United States of America as represented by the Secretary,

Health and Human Services  
Mayer, Ghislaine  
Miller, Louis

<120> PLASMODIUM FALCIPARUM ERYTHROCYTE  
BINDING PROTEIN BAEBL FOR USE AS A VACCINE

<130> NIH209.001PCT

<150> US 60/281130

<151> 2001-04-02

<160> 22

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 4138

<212> DNA

<213> Plasmodium falciparum

<400> 1

tttataatat	cgttttttta	tgagcattat	atcatataaa	taattaatta	aataaatata	60
tatatatata	taattataat	tagaccaata	aattatatat	aatgaaagga	tattttaata	120
tatatTTTTT	aattccttta	atTTTTTTT	ataatgtaat	aagaataaat	gaatcaataa	180
taggtagaac	actttataat	agacaagatg	aatcatcaga	tatttcaagg	gtaaattcac	240
ccgaattaaa	taataatcat	aaaactaata	tatatgattc	agattacgaa	gatgtaaaata	300
ataaattaat	aaacagtttt	gtagaaaata	aaagtgtgaa	aaaaaaaaag	tctttaagtt	360
ttataaataa	taaaacaaaa	tcatatgata	taattccacc	ttcatattca	tataggaatg	420
ataaatttaa	ttcactttcc	gaaaatgaag	ataattctgg	aaatacaaat	agtaataatt	480
tgcgaaatac	ttctgaaata	tctattggaa	aggataataa	acaatatagc	tttatacaga	540
aacgtactca	tttgtttgct	tgtggaataa	aaagaaaatc	aataaaatgg	atatgtcgag	600
aaaacagtga	gaaaattact	gtatgtgttc	ctgatagaaa	aatacaacta	tgtgttgcaa	660
atTTTTTaaa	ctcacgttta	gaaacaatgg	aaaagtttaa	agaaatattt	ttaatttctg	720
ttaatacaga	agcaaaatta	ttatataaca	aaaatgaagg	aaaagatccc	tcaatatatt	780
gtaatgaatt	aagaaatagt	ttttcagatt	ttagaagttc	atTTtataggt	gatgatattg	840
atTTTgtgg	taatacagat	agagtc aaag	gatataTTaa	tacgaagttc	tccgattatt	900
ataaggaaaa	aaatgttgaa	aaattaaata	atatcaaaaa	agaatggtgg	gaaaaaaata	960
aagcaaatTT	gtggaatcac	atgatagtaa	atcataaagg	aaacataagt	aaagaatgtg	1020
ccataattcc	cgcggaagaa	cctcaaatta	atctatggat	aaaagaatgg	aatgaaaact	1080
tcttgatgga	aaagaagaga	ttgtttttta	atataaaaaga	taagtgtgtt	gaaaacaaaa	1140
aatatgaagc	atgttttggt	ggatgtaggc	ttccatgttc	ttcatataca	tcattttatga	1200
aaaaaagtaa	aacacaaatg	gagggttttg	cgaacttgta	taaaaagaaa	aattcaggag	1260
tggataaaaa	taattttctg	aatgatcttt	ttaaaaaaaa	taataaaaaat	gatttagatg	1320
atTTTTTcaa	aaatgaaaag	gaatatgatg	atTTtatgtga	ttgcagatat	actgctacta	1380
ttattaaaag	ttttctaaat	ggtcctgcta	aaaatgatgt	agatatTgca	tcacaaatta	1440
atgttaatga	tcttcgaggg	tttgatgta	attataaaaag	taataatgaa	aaaagttgga	1500
attgtactgg	aacattttacg	aacaaatttc	ctggtacatg	tgaaccccc	agaagacaaa	1560
ctttatgtct	tggacgtaca	tatctttttac	atcgttggtca	tgaggaagat	tataaggaac	1620
atTTactttg	agcttcaata	tatgaggcgc	aattattaaa	atataaaatat	aaggaaaagg	1680
atgaaaatgc	attgtgtagt	ataatacaaa	atagttatgc	agatttgga	gatattatca	1740
aggatcgga	tataataaaa	gattattatg	gtaaaaaaat	ggaagaaaat	ttaaataaag	1800
taaacaaaga	taaaaaacgt	aatgaagaat	ctttgaagat	ttttcgtgaa	aaatggtggg	1860
atgaaaacaa	ggagaatgta	tggaaagtaa	tgtcagcagt	acttaaaaaat	aaggaaacgt	1920
gtaaagatta	tgataagttt	caaaaagattc	ctcaattttt	aagatggttt	aaggaaatggg	1980
gagacgattt	ttgtgagaaa	agaaaagaga	aaatatattc	atttgagtca	tttaaggtag	2040
aatgtaagaa	aaaagattgt	gatgaaaata	catgtaaaaa	taaatgtagt	gaatataaaa	2100

```

aatggataga tttgaaaaaa agtgaatatg agaaacaagt tgataaatac aaaaaagata 2160
aaaataaaaa gatgtatgat aatattgatg aagtaaaaaa taaagaagcc aatgtttact 2220
taaaagaaaa atccaaagaa tgtaaagatg taaatttcga tgataaaatt tttaatgaga 2280
gtccaaatga atatgaagat atgtgtaaaa aatgtgatga aataaaatat ttaaatgaaa 2340
ttaaatatcc taaaacaaaa cacgatatat atgatataga tacattttca gatacttttg 2400
gtgatggaac gccaaataagt attaatgcaa atataaatga acaacaaagt gggaaggata 2460
cctcaaatac tggaaatagt gaaacatcag attcaccggg tagtcatgaa ccagaaagtg 2520
atgctgcaat taatgtagaa aagttaagtg gtgatgaaag ttcaagtga acaagaggaa 2580
tattagatat taatgatcca agtggtacga acaatgtcaa tgaagttcat gatgcttcaa 2640
atacacaagg tagtgtttca aatacttctg atataacgaa tggacattcg gaaagttccc 2700
tgaatagaac aacgaatgca caagatatta aaataggccg ttcaggaaat gaacaaagtg 2760
ataatcaaga aaatagttca cattctagtg ataattcagg ttctttgaca atcggacaag 2820
ttccttcaga ggataatacc caaatacat atgattcaca aaacctcat agagatacac 2880
ctaattgcatt agcatcttta ccatcagatg ataaaattaa tgaatatagag ggtttcgatt 2940
ctagtagaga tagtgaaaat ggtaggggtg atacaacatc aaatactcat gatgtacgtc 3000
gtacgaatat agtaagtgag agacgtgtga atagccatga ttttattaga aacggaatgg 3060
cgaataacaa tgcacatcat caatatataa cgcaaattga gaataatgga atcataagag 3120
gacaagaga aagtgcgggg aatagtgtta attataaaga taatccaaag aggagtaatt 3180
tttctccga aaatgatcat aagaaaaata tacaggaata taattctaga gatactaaaa 3240
gagtaaggga ggaaataatt aaattatcga agcaaaataa atgcaacaat gaatattcca 3300
tggaatatgt tacctattct gacgaaagga atagttcacc gggtccttgt tctagagaag 3360
aaagaaagaa attatgttgt cagatttcag attattgttt aaaatatatt aacttttatt 3420
caattgaaata ttataattgt ataaaatctg aaattaaaag tccagaatat aaatgtttta 3480
aaagcgaggg tcaatcaagt atgtttcata tataatgaaa ataactaaat taaaataaat 3540
atctaaattt tccattttaa tataatttaa ataatttatt tttttattat aaaatttatt 3600
tgtttttttt tatataatgt tttatttttt ctttacaggc attccttatt ttgctgctgg 3660
aggtaattta gttgtaatag tcttactttt gagttcagca tctagaatgg ggaaaagggt 3720
acacgcataa gattaaataa atatgactat actttttaat ttttatatta tttatattat 3780
tattatgttt taacaagaat ataaattata ttatatattt atatgtatat atattttttt 3840
atatagtaat gaagaatatg atataggaga atctaataa gaagcaactt ttgaagaaaa 3900
taattattta aataaactat cgcgcatatg taagattaat ataaaaagtg aaaatttcat 3960
attaaataaa taaatgaata tatatatata tatatatata tatatacatg 4020
ttatataaat ttttaagtta tattatttac ttttttctca gttaatcaag aagtacaaga 4080
gacaaacatt tcagattatt ccgagtacaa ttataatgaa aagaatatgt attaatatt 4138

```

&lt;210&gt; 2

&lt;211&gt; 1210

&lt;212&gt; PRT

&lt;213&gt; Plasmodium falciparum

&lt;400&gt; 2

```

Met Lys Gly Tyr Phe Asn Ile Tyr Phe Leu Ile Pro Leu Ile Phe Leu
1          5          10          15
Tyr Asn Val Ile Arg Ile Asn Glu Ser Ile Ile Gly Arg Thr Leu Tyr
20        25        30
Asn Arg Gln Asp Glu Ser Ser Asp Ile Ser Arg Val Asn Ser Pro Glu
35        40        45
Leu Asn Asn Asn His Lys Thr Asn Ile Tyr Asp Ser Asp Tyr Glu Asp
50        55        60
Val Asn Asn Lys Leu Ile Asn Ser Phe Val Glu Asn Lys Ser Val Lys
65        70        75        80
Lys Lys Arg Ser Leu Ser Phe Ile Asn Asn Lys Thr Lys Ser Tyr Asp
85        90        95
Ile Ile Pro Pro Ser Tyr Ser Tyr Arg Asn Asp Lys Phe Asn Ser Leu
100       105       110
Ser Glu Asn Glu Asp Asn Ser Gly Asn Thr Asn Ser Asn Asn Phe Ala
115       120       125
Asn Thr Ser Glu Ile Ser Ile Gly Lys Asp Asn Lys Gln Tyr Thr Phe
130       135       140
Ile Gln Lys Arg Thr His Leu Phe Ala Cys Gly Ile Lys Arg Lys Ser
145       150       155       160
Ile Lys Trp Ile Cys Arg Glu Asn Ser Glu Lys Ile Thr Val Cys Val
165       170       175

```

Pro	Asp	Arg	Lys	Ile	Gln	Leu	Cys	Val	Ala	Asn	Phe	Leu	Asn	Ser	Arg
			180					185					190		
Leu	Glu	Thr	Met	Glu	Lys	Phe	Lys	Glu	Ile	Phe	Leu	Ile	Ser	Val	Asn
		195					200					205			
Thr	Glu	Ala	Lys	Leu	Leu	Tyr	Asn	Lys	Asn	Glu	Gly	Lys	Asp	Pro	Ser
	210					215					220				
Ile	Phe	Cys	Asn	Glu	Leu	Arg	Asn	Ser	Phe	Ser	Asp	Phe	Arg	Ser	Ser
225				230						235					240
Phe	Ile	Gly	Asp	Asp	Met	Asp	Phe	Gly	Gly	Asn	Thr	Asp	Arg	Val	Lys
			245						250					255	
Gly	Tyr	Ile	Asn	Thr	Lys	Phe	Ser	Asp	Tyr	Tyr	Lys	Glu	Lys	Asn	Val
			260					265						270	
Glu	Lys	Leu	Asn	Asn	Ile	Lys	Lys	Glu	Trp	Trp	Glu	Lys	Asn	Lys	Ala
		275					280					285			
Asn	Leu	Trp	Asn	His	Met	Ile	Val	Asn	His	Lys	Gly	Asn	Ile	Ser	Lys
		290				295					300				
Glu	Cys	Ala	Ile	Ile	Pro	Ala	Glu	Glu	Pro	Gln	Ile	Asn	Leu	Trp	Ile
305					310					315					320
Lys	Glu	Trp	Asn	Glu	Asn	Phe	Leu	Met	Glu	Lys	Lys	Arg	Leu	Phe	Leu
			325						330					335	
Asn	Ile	Lys	Asp	Lys	Cys	Val	Glu	Asn	Lys	Lys	Tyr	Glu	Ala	Cys	Phe
			340					345					350		
Gly	Gly	Cys	Arg	Leu	Pro	Cys	Ser	Ser	Tyr	Thr	Ser	Phe	Met	Lys	Lys
		355					360					365			
Ser	Lys	Thr	Gln	Met	Glu	Val	Leu	Thr	Asn	Leu	Tyr	Lys	Lys	Lys	Asn
		370				375					380				
Ser	Gly	Val	Asp	Lys	Asn	Phe	Leu	Asn	Asp	Leu	Phe	Lys	Lys	Lys	Asn
385				390					395						400
Asn	Lys	Asn	Asp	Leu	Asp	Asp	Phe	Phe	Lys	Asn	Glu	Lys	Glu	Tyr	Asp
			405						410					415	
Asp	Leu	Cys	Asp	Cys	Arg	Tyr	Thr	Ala	Thr	Ile	Ile	Lys	Ser	Phe	Leu
			420					425					430		
Asn	Gly	Pro	Ala	Lys	Asn	Asp	Val	Asp	Ile	Ala	Ser	Gln	Ile	Asn	Val
		435					440					445			
Asn	Asp	Leu	Arg	Gly	Phe	Gly	Cys	Asn	Tyr	Lys	Ser	Asn	Asn	Glu	Lys
		450				455					460				
Ser	Trp	Asn	Cys	Thr	Gly	Thr	Phe	Thr	Asn	Lys	Phe	Pro	Gly	Thr	Cys
465				470						475					480
Glu	Pro	Pro	Arg	Arg	Gln	Thr	Leu	Cys	Leu	Gly	Arg	Thr	Tyr	Leu	Leu
			485						490					495	
His	Arg	Gly	His	Glu	Glu	Asp	Tyr	Lys	Glu	His	Leu	Leu	Gly	Ala	Ser
			500					505					510		
Ile	Tyr	Glu	Ala	Gln	Leu	Leu	Lys	Tyr	Lys	Tyr	Lys	Glu	Lys	Asp	Glu
		515					520					525			
Asn	Ala	Leu	Cys	Ser	Ile	Ile	Gln	Asn	Ser	Tyr	Ala	Asp	Leu	Ala	Asp
		530				535					540				
Ile	Ile	Lys	Gly	Ser	Asp	Ile	Ile	Lys	Asp	Tyr	Tyr	Gly	Lys	Lys	Met
545				550						555					560
Glu	Glu	Asn	Leu	Asn	Lys	Val	Asn	Lys	Asp	Lys	Lys	Arg	Asn	Glu	Glu
			565						570					575	
Ser	Leu	Lys	Ile	Phe	Arg	Glu	Lys	Trp	Trp	Asp	Glu	Asn	Lys	Glu	Asn
			580					585					590		
Val	Trp	Lys	Val	Met	Ser	Ala	Val	Leu	Lys	Asn	Lys	Glu	Thr	Cys	Lys
		595					600					605			
Asp	Tyr	Asp	Lys	Phe	Gln	Lys	Ile	Pro	Gln	Phe	Leu	Arg	Trp	Phe	Lys
		610				615					620				
Glu	Trp	Gly	Asp	Asp	Phe	Cys	Glu	Lys	Arg	Lys	Glu	Lys	Ile	Tyr	Ser
625				630						635					640
Phe	Glu	Ser	Phe	Lys	Val	Glu	Cys	Lys	Lys	Lys	Asp	Cys	Asp	Glu	Asn
			645						650					655	
Thr	Cys	Lys	Asn	Lys	Cys	Ser	Glu	Tyr	Lys	Lys	Trp	Ile	Asp	Leu	Lys
			660					665					670		
Lys	Ser	Glu	Tyr	Glu	Lys	Gln	Val	Asp	Lys	Tyr	Thr	Lys	Asp	Lys	Asn

Page 4

Arg Ile Phe Asn Gln Glu Val Gln Glu Thr Asn Ile Ser Asp Tyr Ser  
 1185 1190 1195 1200  
 Glu Tyr Asn Tyr Asn Glu Lys Asn Met Tyr  
 1205 1210

<210> 3  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> primer

<400> 3  
 agaccaataa attatatata atgaaagga 29

<210> 4  
 <211> 27  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> primer

<400> 4  
 tttaaacttt tccattgttt ctaaacg 27

<210> 5  
 <211> 28  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> primer

<400> 5  
 ataaatttaa ttcactttcc gaaaatga 28

<210> 6  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> primer

<400> 6  
 aaaacaatct cttcttttcc atcaag 26

<210> 7  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> primer

<400> 7  
 tttataggtg atgatatgga ttttgg 26

<210> 8  
 <211> 25

<212> DNA  
<213> Artificial Sequence

<220>  
<223> primer

<400> 8  
tcgtaaatgt tccagtacaa ttcct 25

<210> 9  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> primer

<400> 9  
caaatggagg ttttgacgaa cttg 24

<210> 10  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> primer

<400> 10  
taagtactgc tgacattact ttcca 25

<210> 11  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> primer

<400> 11  
ggagcttcaa tatatgaggc gca 23

<210> 12  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> primer

<400> 12  
atatcttcat attcatttgg actctc 26

<210> 13  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> primer

<400> 13  
tgagtcattt aaggtagaat gtaaga 26

<210> 14  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> primer

<400> 14  
ggaactttcc gaatgtccat tcgt

24

<210> 15  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> primer

<400> 15  
taaatgaaca acaaagtggg aagga

25

<210> 16  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> primer

<400> 16  
attctcaatt tgcgttatat attgatg

27

<210> 17  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> primer

<400> 17  
agttccttca gaggataata ccca

24

<210> 18  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> primer

<400> 18  
cttgattgac cctcgctttt aaaac

25

<210> 19  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> primer



<400> 19  
actaaaagag taaggaggga aataat 26

<210> 20  
<211> 31  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> primer

<400> 20  
tataaaatac attgaattat ttaaactatt g 31

<210> 21  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 21  
attccttatt ttgctgctgg aggt 24

<210> 22  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 22  
aagttgcttc tatattagat tctcct 26